


DESI Technical Design Report Part I: Science, Targeting, and Survey Design

July 15, 2015

Rev A



DOE CD-2 Independent Review Panel Version
Rev A incorporating committee comments
2015-07-27

Compiled: 2015-08-27 15:40
DESI-1125 Part I

Contents

1	Overview of DESI	1
2	Science Motivation and Requirements	3
2.1	Introduction	3
2.2	Measuring Distances with Baryon Acoustic Oscillations	7
2.2.1	Theory	7
2.2.2	BAO in Galaxies	7
2.2.2.1	Observational Systematics	9
2.2.2.2	Theoretical Systematics	9
2.2.3	BAO in the Ly- α Forest	10
2.2.3.1	Systematics	11
2.3	Measuring Growth of Structure with Redshift Space Distortions	13
2.3.1	Theory	13
2.3.2	Systematics	14
2.3.3	Current Status of RSD Measurements	15
2.4	Distance, Growth, Dark Energy, and Curvature Constraint Forecasts	18
2.4.1	Forecasting Overview	18
2.4.2	Baseline Survey	20
2.4.3	Summary of Forecasts	23
2.4.4	Forecasting Details	29
2.4.4.1	Galaxy and Quasar Clustering	29
2.4.4.2	Ly- α Forest	29
2.5	Cosmology Beyond Dark Energy	31
2.5.1	Inflation	31
2.5.1.1	Spectral Index and Its Running	31
2.5.1.2	Primordial non-Gaussianity	32
2.5.2	Neutrinos	35
2.5.2.1	Neutrino Mass	35
2.5.2.2	Dark Radiation (e.g., sterile neutrinos)	36
2.6	The Milky Way Survey: Near-Field Cosmology from Stellar Spectroscopy	38
2.7	Complementarity with Other Surveys	39
2.7.1	Synergies with <i>Planck</i> and Future CMB Experiments	39
2.7.2	Synergies of DESI with DES and LSST	39
2.7.3	Synergies of DESI with <i>Euclid</i> / <i>WFIRST</i>	41
3	Target Selection	43
3.1	Targets: Bright Galaxy Sample	45
3.1.1	Overview of the Sample	45
3.1.2	Sample Properties	46
3.2	Targets: Luminous Red Galaxies	48
3.2.1	Overview of the Sample	48
3.2.2	Selection Technique for $z > 0.6$ LRGs	49
3.2.3	Sample Properties	50
3.3	Targets: Emission Line Galaxies	54
3.3.1	Overview of the sample	54

3.3.2	Selection Technique for $z > 0.6$ ELGs	55
3.3.3	Sample Properties	56
3.4	Targets: QSOs	60
3.4.1	Overview of the sample	60
3.4.2	Selection Technique	61
3.4.3	Sample Properties	63
3.4.4	Near- and mid-term developments for QSO target selection	65
3.4.5	Variability Data Improves Selection of High-Redshift QSOs	65
3.5	Calibration Targets	66
3.6	Baseline Imaging Datasets	67
3.6.1	Blanco/DECam Surveys ($\text{DEC} \leq 30^\circ$)	67
3.6.2	Bok/90Prime Survey ($\text{DEC} \geq 30^\circ$)	69
3.6.3	Mayall/MOSAIC Survey ($\text{DEC} \geq 30^\circ$)	70
3.6.4	<i>WISE</i> All-Sky Survey	71
3.7	Additional Imaging Data	71
3.7.1	SDSS	72
3.7.2	PanSTARRS-1	72
3.7.3	PTF, iPTF, and ZTF	72
3.7.4	CFHT	73
3.7.5	SCUSS	73
3.8	The Tractor Photometry for Target Selection	73
4	Survey Design	76
4.1	Introduction	76
4.2	Survey Footprint	76
4.3	Field Centers	76
4.4	Observation Strategy	78
4.4.1	Sequence of Observations	78
4.4.2	Exposure Times and Margin	80
4.5	The Bright Galaxy and Milky Way Surveys	82
4.5.1	Introduction	82
4.5.2	Survey Footprint	82
4.5.3	Field Centers	82
4.5.4	Observation Strategy	83
A	Acronyms	84
	References	90

1 Overview of DESI

DESI is a Stage IV ground-based dark energy experiment that will study baryon acoustic oscillations (BAO) and the growth of structure through redshift-space distortions (RSD) with a wide-area galaxy and quasar redshift survey. DESI is the successor to the successful Stage-III BOSS redshift survey and complements imaging surveys such as the Stage-III Dark Energy Survey (DES, operating 2013–2018) and the Stage-IV Large Synoptic Survey Telescope (LSST, planned start early in the next decade). DESI is an important component of the DOE Cosmic Frontier program, meeting the need for a wide-field spectroscopic survey identified in the 2011 “Rocky-III” dark energy community planning report. In addition to providing Stage IV constraints on dark energy, DESI will provide new measurements that can constrain theories of modified gravity and inflation, and that will measure the sum of neutrino masses.

The DESI instrument is a robotically-actuated, fiber-fed spectrograph capable of taking up to 5,000 simultaneous spectra over a wavelength range from 360 nm to 980 nm. The fibers feed ten three-arm spectrographs with resolution $R = \lambda/\Delta\lambda$ between 2000 and 5500, depending on wavelength. This powerful instrument will be installed at prime focus on the 4-m Mayall telescope in Kitt Peak, Arizona, along with a new optical corrector, which will provide a three-degree diameter field of view. The DESI collaboration will also deliver a spectroscopic pipeline and data management system to reduce and archive all data for eventual public use.

The DESI instrument will be used to conduct a five-year survey designed to cover 14,000 deg². To trace the underlying dark matter distribution, spectroscopic targets will be selected in four classes from imaging data. We will measure luminous red galaxies (LRGs) up to $z = 1.0$, extending the BOSS LRG survey in both redshift and survey area. To probe the Universe out to even higher redshift, DESI will target bright [O II] emission line galaxies (ELGs) up to $z = 1.7$. Quasars will be targeted both as direct tracers of the underlying dark matter distribution and, at higher redshifts ($2.1 < z < 3.5$), for the Ly- α forest absorption features in their spectra, which will be used to trace the distribution of neutral hydrogen. When moonlight prevents efficient observations of the faint targets of the baseline survey, DESI will conduct a magnitude-limited Bright Galaxy Survey (BGS) comprising approximately 10 million galaxies with a median $z \approx 0.2$. In total, more than 30 million galaxy and quasar redshifts will be obtained to measure the BAO feature and determine the matter power spectrum, including redshift space distortions.

In the following document, we primarily refer to this baseline survey, which would span 14,000 deg². We also calculate numbers for a minimum survey spanning 9,000 deg², which is still sufficient to meet the requirements of a Stage-IV project.

DESI provides at least an order of magnitude improvement over BOSS both in the comoving volume it probes and the number of galaxies it will map. This will significantly advance our understanding of the expansion history of the Universe, providing more than thirty sub-percent-accuracy distance measurements. Precision on the expansion history of the Universe is a powerful probe of the nature of dark energy. This can be quantified with the Dark Energy Task Force figure of merit (DETF FoM), which measures the combined precision on the dark energy equation of state today, w_0 , and its evolution with redshift w_a . DESI galaxy BAO measurements achieve a DETF FoM of 133, more than a factor of three better than the DETF FoM of all Stage-III galaxy BAO measurements combined. The FoM increases to 169 with the inclusion of Ly- α forest BAO, and 332 including galaxy broadband power spectrum to $k = 0.1 \, h \, \text{Mpc}^{-1}$. DESI clearly satisfies the DETF criteria for a Stage-IV experiment. Moreover, the FoM grows to 704 when the galaxy broadband power spectrum data out to $k < 0.2 \, h \, \text{Mpc}^{-1}$ are included.

In addition, DESI will measure the sum of neutrino masses with an uncertainty of 0.020 eV (for $k_{\text{max}} < 0.2 \, h \, \text{Mpc}^{-1}$), sufficient to make the first direct detection of the sum of the neutrino masses

at $3\text{-}\sigma$ significance and rule out the the inverted mass hierarchy at 99% CL, if the hierarchy is normal and the masses are minimal. DESI will also place significant constraints on theories of modified gravity and of inflation by measuring the spectral index n_s and its running with wavenumber, α_s . The BGS will enable the best ever measurements of low redshift BAO and RSD, including the use of multiple-tracer methods that exploit galaxy populations with different clustering properties, and it will yield novel tests of modified gravity theories using the velocity fields of cluster infall regions. Because the nearby galaxies of the BGS are too clustered to fill all of the targets, in parallel with the BGS, DESI will conduct a survey of Milky Way stars, that can be used to trace the dark matter halo of the Milky Way and probe the small-scale structure of Λ CDM.

DESI will provide an unprecedented multi-object spectroscopic capability for the U.S. through an existing NSF telescope facility. Many other science objectives can be addressed with the DESI wide field survey dataset and through bright time and piggy-back observation programs. Much as with SDSS, a rich variety of projects will flow from the legacy data from the DESI survey.

DESI will overlap with the DES and LSST survey areas, which are primarily in the Southern hemisphere but which will have equatorial and northern ecliptic regions. DESI will be a pathfinder instrument for the massive spectroscopic follow-up required for future large area imaging surveys such as LSST.

This portion of the Technical Design Report summarizes the DESI scientific goals, the target selection, and survey design. The accompanying portion of the TDR describes the instrument and optical design, integration and test plan, and the data management system. The companion Science Requirements Document provides information that guides the design. The DESI construction management plan is presented in the accompanying Project Execution Plan. Likewise, project cost and schedule are available in appropriate Project Office documents.

2 Science Motivation and Requirements

2.1 Introduction

DESI will explore some of the most fundamental questions in physical science: what is the composition of the Universe at large and what is the nature of space-time? These questions are now open to exploration because of recent discoveries. We summarize here the framework used to express these questions and the parameters used to quantify our understanding.

There are several pillars of the cosmological model that are now well established: 1) a period of rapid acceleration — *inflation* or a similar process — occurred in the early Universe, generating the primordial fluctuations, which seeded large scale structures, galaxies and galaxy clusters, which grew during the decelerating, matter dominated era 2) gravitational instabilities produced acoustic oscillations in the plasma, which were imprinted about 400,000 years after this inflation period, when photons decoupled from atoms and produced the Cosmic Microwave Background 3) this was followed by a period of matter domination, when small density fluctuations grew into large-scale structure, 4) comparatively recently, there was a transition to accelerated expansion driven by either a modification to General Relativity or a new form of energy — *dark energy* — not due to any particles known or unknown, and which contributes about 68% of the Universe's energy density, and 5) about four-fifths of the 27% of the energy density today is due to matter outside the Standard Model of particle physics — *dark matter* — which is responsible for large-scale structure formation and accounts for galaxy rotation curves and the motions of galaxies in clusters.

That the Universe is expanding more and more rapidly was first revealed through measurements of Type Ia supernovae [1, 2], and subsequently confirmed using other techniques. Within General Relativity, accelerated expansion requires $\rho + 3p < 0$, where ρ is the total energy density and p is the total pressure of the matter, radiation, and other ingredients. The total equation of state $w = p/\rho$ must be less than $-1/3$ for accelerated expansion. The equation of state need not be a constant; in general it depends on time, or equivalently the scale size of the universe $a = 1/(1+z)$. From now on, we let w denote the equation of state of the dark energy component alone.

For ordinary non-relativistic matter, the pressure is negligible compared to the energy due to the rest mass and thus $w = 0$. For photons and other massless particles, $w = 1/3$. The cosmological constant term is equivalent to dark energy with $w = -1$. Generally, energy with an equation of state $w(a)$ evolves as $\rho(a) = \rho(a=1)F(a)$, where $F(a) = 1$ for a cosmological constant and for a general equation of state $w(a)$ is

$$F(a) \equiv \exp \left[3 \int_a^1 \frac{da'}{a'} (1 + w(a')) \right]. \quad (2.1)$$

The contributions to the energy density of the Universe are conventionally expressed relative to the critical density

$$\rho_{\text{crit}} = \frac{3H_0^2}{8\pi G}, \quad (2.2)$$

which would be just sufficient to slow the expansion ultimately to zero in the absence of a dark energy component Ω_Λ . We write

$$\Omega_m = \frac{\rho_m}{\rho_{\text{crit}}}. \quad (2.3)$$

We define Ω_r for radiation and Ω_{DE} for dark energy analogously. The curvature term $\Omega_k = -k/H_0^2$ is defined so that General Relativity requires for a Universe with spatial curvature k

$$\Omega_r + \Omega_m + \Omega_k + \Omega_{DE} = 1. \quad (2.4)$$

The expansion rate of the Universe is given by

$$H(a) \equiv \frac{\dot{a}}{a} = H_0 [\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_{DE} F(a)]^{1/2}. \quad (2.5)$$

The contribution from radiation, Ω_r is negligible today and inflation predicts that the curvature is zero. The Hubble constant today is $H_0 = h \times 100 \text{ km/s/Mpc} \approx 70 \text{ km/s/Mpc}$.

It is standard to parameterize the equation of state as

$$w(a) = w_0 + (1 - a)w_a, \quad (2.6)$$

which accurately reproduces distances for a wide range of models.

We have three possible explanations for the accelerating expansion of the Universe: a cosmological constant, equivalent to static dark energy with $w = -1$; a dynamical dark energy with $w(a) \neq -1$; or a failure of General Relativity. DESI is designed to address this fundamental question about the nature of the Universe. The challenge of distinguishing the cosmological constant solution from dark energy with w near -1 is displayed in Figure 2.1.

The Dark Energy Spectroscopic Instrument (DESI) [3] will provide precise spectroscopic redshifts of more than thirty million objects. From these will come three-dimensional maps of the distribution of matter covering unprecedented volume. DESI will survey an enormous volume at $0.4 < z < 3.5$ using luminous red galaxies, emission line galaxies, and quasars, producing tight constraints on the large-scale clustering of the Universe. In addition, DESI will perform a Bright Galaxy Survey (BGS) of the $z < 0.4$ Universe, allowing the study of cosmic structure in the dark-energy-dominated epoch with much denser sampling. These data will help establish whether cosmic acceleration is due to a mysterious component of the Universe or a cosmic-scale modification of GR, and will constrain models of primordial inflation.

DESI will have a dramatic impact on our understanding of dark energy through its primary measurement, that of baryon acoustic oscillations. Waves that propagated in the electron-photon-baryon plasma before recombination imprint a feature at a known comoving physical scale (150 Mpc or $4.6 \times 10^{24} \text{ m}$) in the distribution of separations between pairs of galaxies. Localizing this baryon acoustic oscillation (BAO) feature and comparing its apparent size to the known physical scale provides a measurement of the distance to the galaxy sample and thus the expansion history of the Universe. The BAO measurement was singled out by the Dark Energy Task Force [4] as having the fewest experimental uncertainties among the techniques for measuring dark energy; it simply depends on the galaxy locations, rather than their shapes or brightnesses. DESI's two-point correlation measurements will also detect the anisotropies in galaxy clustering – redshift space distortions (RSD) – due to the peculiar velocities of galaxies generated by density perturbations. This gives a direct measurement of the properties of gravity at each redshift, through its effect on galaxies' motions.

In addition to the constraints on dark energy, the galaxy and Ly- α flux power spectra will reflect signatures of neutrino mass, scale dependence of the primordial density fluctuations from inflation, and possible indications of modified gravity. To realize the potential of these techniques requires an enormous number of redshifts over a deep, wide volume and thus a substantial investment in a new, more capable instrument and significant allocation of the telescope time.

The DESI survey will have considerable impact beyond these cosmological highlights on the study of galaxies, quasars, and stars. Spectroscopy is a core tool of astrophysics, and the ability to combine many millions of spectra with modern wide-field, multi-wavelength imaging surveys will yield rich opportunities. While the DESI collaboration includes members planning to work on these topics, we do not discuss these in this design report, as they are not driving requirements. We




Figure 2.1: The expansion history of the Universe for different models of dark energy, holding the present-day Hubble constant fixed. The inset shows the spacing between five models with constant w ranging from -0.97 to -1.03 , showing the exquisite precision required to distinguish these. Overlaid are measurements of the distance-redshift relation, translated into errors on lookback time at each redshift. Measurements from current supernovae, binned in redshift, are shown in blue; current BAO measurements from BOSS DR9, WiggleZ, and 6dF are shown in red; projections for DESI are shown in black. DESI measurements have the ability to make very tight constraints on dark energy, although we caution that this figure shows variations in only one cosmological parameter. Full forecasts, such as those presented in § 2.4.3, must marginalize over other cosmological parameters such as Ω_m and H_0 .

make one brief exception for the Milky Way Survey (§ 2.6), as it will involve a substantial number of targets that piggyback on the Bright Galaxy Survey, using fibers that have no suitable galaxy available within their patrol radius.

2.2 Measuring Distances with Baryon Acoustic Oscillations

DESI will measure the expansion of the Universe by observing the imprint of baryon acoustic oscillations set down in the first 380,000 years of its existence. This pattern has the same source as the pattern seen in the cosmic microwave background, but DESI will map it as a function of cosmic time, while the CMB can see it only at one instant. The pattern is imprinted on all matter at large scales and can be viewed by observing galaxies of various kinds or by observing the distribution of neutral hydrogen across the cosmos, and shows up as excess correlations at the characteristic distance of the sound horizon at decoupling.

2.2.1 Theory

Initial fluctuations in density and pressure provided sources for sound waves that propagated in the photon-electron-baryon plasma of the early Universe (see, for example, [5]). These sound waves propagated with a speed approximately $c/\sqrt{3}$ until the Universe cooled sufficiently for electrons and ions to recombine to neutral atoms, causing the sound speed to drop dramatically. An excess of matter was left both at the source of the wave and at the surface where these waves terminated. The matter excesses at these locations left their imprint on the large-scale structure of galaxies and hydrogen gas. Before a wave stopped, it traveled a co-moving distance $s \approx 150$ Mpc, which can be computed to precision 0.3% from cosmological parameters extremely well measured in CMB.

Viewed transversely, the 150-Mpc ruler subtends an angle θ such that

$$s = (1+z)D_A(z)\theta = \theta \int_0^z \frac{c dz'}{H(z')} \quad (2.7)$$

where $D_A(z)$ is the angular-diameter distance to an object at redshift z . The final equality holds only if the curvature is zero.

While the CMB gives us a purely angular correlation function, the characteristic scale is present in the three-dimensional distribution of large-scale structure. Viewed along the line of sight, correlations are enhanced for galaxy pairs separated by Δz such that

$$\frac{c\Delta z}{H(z)} \approx s \quad (2.8)$$

This latter measurement requires a spectroscopic survey to resolve the full three-dimensional density distribution of galaxies.

The observation of the peak in the two-point correlation function thus provides a means of measuring both the angular diameter distance, $D_A(z)$ and the Hubble expansion rate, $H(z)$. The ability of the BAO method to directly probe $H(z)$ is unique among dark energy probes. This becomes progressively more important at higher redshifts since H measures the instantaneous expansion rate (and through it, the total energy density of the Universe) while D_A measures the integrated expansion history. Measuring both improves our ability to distinguish between different cosmological models.

2.2.2 BAO in Galaxies

The best developed application of the BAO technique uses galaxies as tracers of the matter distribution; the BAO feature appears in the two-point correlation function of galaxies, the probability, in excess of random, that two galaxies are separated by a distance r . This has been achieved with high statistical significance in several measurements spanning the redshift range from $z = 0$ to $z = 1$.




Figure 2.2: The angle-averaged correlation functions [left] and power spectra [right], before [top] and after [bottom] reconstruction measured using the BOSS DR11 CMASS galaxy sample [6]. The BAO feature is clearly detected at over 7σ as a peak in the correlation function and a corresponding set of oscillations in the power spectrum.

The highest significance detection ($> 7\sigma$) is currently that of the Baryon Oscillation Spectroscopic Survey (BOSS) using the $z > 0.45$ sample [6]. We show representative data in Figure 2.2. These data measured the distance-like quantity $D_V(z) \equiv ((1+z)D_A)^{2/3}(cz/H(z))^{1/3}$ to a redshift of 1.0%, the most precise measurement using the BAO technique. The lower redshift $z < 0.45$ sample in BOSS constrained the same combination of distances to 2%. At still lower redshifts, the 6-Degree Field Galaxy Redshift Survey [7] measured the distance to $z = 0.106$ with 4.5% accuracy. At a somewhat higher redshift, the WiggleZ galaxy survey measured the distance to a redshift of 0.7 to 4% [8]. This combination of these measurements has for the first time enabled mapping the distance-redshift relation purely from BAO measurements.

Most of these measurements used the galaxy correlation function averaged over the orientation of the pair to the line of sight to measure D_V , a combination of D_A and H . More recent work has also measured the correlation functions transverse and parallel to the line of sight, allowing one to break the degeneracy between D_A and H that exists in purely angle averaged measurements.

The current generation of surveys is an excellent proving ground for analysis techniques. For instance, the BOSS experiment compared analyses done in Fourier and in configuration space and used different algorithms for estimating distances from the resulting two-point functions. All these yielded consistent distance measurements, given the statistical precision of the measurements. While the level of consistency is not at the level required by DESI, ongoing surveys provide a clear roadmap for developing and validating improvements to these analysis techniques. The current measurements provide an important validation of our forecasts for DESI presented below.

The non-linear evolution of the matter density field broadens the acoustic peak, potentially decreasing the precision on the distance measurement, and causes a small shift in the peak location, thereby biasing the distance. Ref. [9] pointed out that because this broadening is caused by the large-scale velocity flows resulting from gravitational forces, the effect may be substantially reversed by estimating the velocity fields from the large-scale structure map and moving the galaxies back

to their initial positions. In addition to a notable improvement in the recovered statistical errors, this reconstruction also mitigates the shifts in the distance scale due to nonlinear evolution, with numerical tests showing suppression to below 0.1%. Reconstruction was first applied to the SDSS-II galaxy survey [10], improving the statistical precision by a factor of 1.7. Galaxy samples from latest SDSS data release, DR11, yield similar improvements after reconstruction. See Fig. 2.2. As with the other analysis methods, we expect improvements to reconstruction algorithms before the DESI measurements become available. We however choose to be conservative and assume a reconstruction performance similar to what has already been demonstrated with current data.

2.2.2.1 Observational Systematics

The BAO method is simple in principle — all one requires are the three-dimensional positions of galaxies. The need to preserve the BAO feature along the line of sight sets the requirement on redshift precision. This precision, as stated in the Level 2 Survey Data Set Requirements is $\sigma_z/(1+z) \sim 0.0005$ per galaxy, which is easily within the state-of-the-art and which is achieved throughout our wavelength range in the spectrograph design.

The angular and radial selection functions of the survey can induce systematic uncertainties. The angular selection function is determined by the imaging survey used for targeting, and may be spuriously modulated by photometric calibrations, seeing and extinction variations, and image deblending. All of these effects are intrinsically angular effects and therefore may be separated from the BAO feature which is a feature in three-dimensional physical space (not isolated to the angular degrees of freedom). A similar separation is possible for systematics in the radial selection function of the survey. The impact of these is therefore expected to be small. In addition, there has been considerable work [11, 12] developing techniques to further mitigate these effects.

The ongoing BAO surveys provide the opportunity to identify and quantify observational systematics. DESI will benefit greatly from this work, but it also faces some unique challenges. The most important of these arise from the fiber positioning system and from the forest of sky lines, which impinge on the radial selection function. The limited patrol radius of the fiber positioners causes the highest density regions to be sampled less completely than lower density regions. This particularly affects the observer’s line of sight and can skew the anisotropic correlation pattern. High sky brightness at certain wavelengths makes it hard to find [OII] emission lines, thereby dropping the spectroscopic completeness at specific redshifts. Initial studies have shown that these survey artifacts can influence the measured clustering, but we expect both to be correctable to good accuracy, as the source of the variations can be tracked with high fidelity. Finding the optimal method to achieve the full statistical precision inherent in the data is an ongoing project of the science team.

2.2.2.2 Theoretical Systematics

The robustness and accuracy of the BAO method derive from the simplicity of the early Universe and the precision with which we know the speed and time of propagation of sound waves in the primordial plasma. The evolution of density fluctuations in the Universe is very well described by linear perturbation theory and is now exquisitely tested by the recent measurements of temperature fluctuations in the Cosmic Microwave Background radiation by the *Planck* satellite [13]. The current CMB measurements constrain the size of the BAO standard ruler to 0.3%. This uncertainty is folded into our forecasts for DESI. Furthermore, any miscalibrations in the acoustic scale would affect principally the determination of the Hubble constant, not the dark energy constraints [14].

The sound waves travel a comoving distance of 150 Mpc, setting the BAO scale to be much larger

than the scale of gravitational collapse even in the present Universe (about 10 Mpc). Analytical calculations, verified by direct numerical simulations, have found the nonlinear evolution of the density field alters the BAO scale by 0.3% at the present epoch, and even less at the higher redshifts probed by DESI.

Galaxy formation may result in an additional shift in the BAO scale due to mismatched weighting of high and low density regions. Initial perturbative and numerical work [15, 16, 17, 18, 19, 20, 21] also find these shifts to be small, with the most extreme shifts of order 0.5%. As discussed above, density-field reconstruction applied to simulations reduces these shifts to the 0.1% level without the need for further modeling. We expect that further modeling from theory and simulations will allow us to robustly limit these uncertainties to well below the DESI statistical limits. In addition, the DESI target samples are designed to overlap in multiple redshift ranges, allowing empirical tests of the robustness of the BAO measurements to different tracer populations.

A recently discovered astrophysical effect that could affect the BAO feature arises from the relative velocities of the baryons and the dark matter at the recombination epoch [22]. This modulates the formation of the earliest protogalaxies and potentially could persist to their descendants (some of which would be measured by DESI). This modulation is due to the same pressure forces that create the BAO, and the impact could shift the measured acoustic scale. This effect is expected to be negligible for the galaxies probed by DESI. Furthermore, [23] demonstrate that this effect would also create a distinctive three-point function signal measurable in DESI that would diagnose any contamination from this effect.

All of the above strongly argue that the theoretical systematic effects associated with the BAO-scale measurements are either intrinsically or correctable to below the 0.1% level required by DESI.

2.2.3 BAO in the Ly- α Forest

Measuring BAO with galaxies as tracers is a mature method [24, 8]. Such measurements become much more difficult for $z \gtrsim 2.0$ where galaxy redshifts are harder to get. However, measuring dark energy properties at this high redshift allows us to probe the Universe well before the advent of accelerated expansion. An interesting possibility is that dark energy density does not become completely negligible at high redshift, as predicted by the cosmological constant or other models with $w \simeq -1$, but rather remains at a level predicted by some particle-physics models and detectable by future surveys [25, 26, 27, 28, 29, 30]. Such a component can only be measured or excluded by a technique sensitive to the expansion history at high redshift.

The Ly- α forest provides the means to measure BAO at redshifts larger than 2. The forest is a collection of absorption features in the spectra of distant quasars blue-ward of the Ly- α emission line [31]. These features arise because the light from a quasar is absorbed by neutral hydrogen in the intergalactic medium. Since the quasar light is constantly red-shifting, hydrogen at different redshifts absorbs at different observed wavelengths in the quasar spectrum. The amount of absorption reflects the local density of neutral hydrogen, which in turn traces the dark matter field on sufficiently large scales.

Numerical simulations and analytical work show that for plausible scenarios, the Ly- α forest is well within the linear biasing regime of scales relevant for BAO [32, 33, 34]. Therefore, measuring three-dimensional correlations in the flux fluctuations of the Ly- α forest provides an accurate method for detecting BAO correlations [32, 35, 36, 37]

Using the Ly- α forest to measure the three-dimensional structure of the Universe became possible with the advent of BOSS, which was the first survey to have a sufficiently high density of quasars to measure correlations on truly cosmological scales. This was done in 2011 [38]. At the beginning of 2013, the first detection of BAO in the Ly- α forest was published in a series of papers




Figure 2.3: Correlation functions of Ly- α forest flux fluctuations based on the BOSS DR11 quasars [42], binned in the cosine of the angle to the line of sight, μ ($\mu = 1$ is along the line of sight, $\mu = 0$ is perpendicular to the line of sight). From left to right, the bins are $\mu > 0.8$, $0.5 < \mu < 0.8$ and $\mu < 0.5$. The points are the measured correlation function, the solid line is the best fit model, while the dashed line is the best fit assuming a fiducial cosmology. These results measure the optimal combination $D_A^{0.3} H^{-0.7}$ to 2%.

[39, 40, 41]. These were recently updated to the almost complete BOSS sample in [42] (Fig. 2.3) yielding a 5σ detection of the BAO feature.

The redshift-space distortions in the Ly- α forest are larger than in galaxy-based measurements [38, 32]. Thus the signal-to-noise for the radial modes is considerably higher than for transverse modes. Consequently, in contrast to the galaxy measurements, the Ly- α forest BAO measurements measure the Hubble parameter $H(z)$ with greater precision than the angular diameter distance $D_A(z)$. For instance, [42] find that the combination $D_A^{0.3} H^{-0.7}$ is optimally constrained to $\sim 2\%$.

2.2.3.1 Systematics

Inevitably, there will be systematic effects that could distort the Ly- α measurements, but these should produce broadband contamination and would not affect our ability to measure an isolated feature in the data, such as the BAO peak. However, unless carefully accounted for, these systematics could contaminate secondary science, such as Ly- α broadband power measurements, neutrino masses and warm dark matter constraints.

Astrophysical contaminants include sources of non-gravitational large scale fluctuations, such as He II reionization and fluctuations in the photo-ionization background. [43, 44, 45, 46, 47] There are also targeting systematics – quasars with significant absorption in the forest region are considerably easier to target, since they are easier to distinguish from stars. As a result, observed Ly- α forest regions are not sampling the Universe randomly, but prefer overdense lines of sight. Back-of-the-envelope calculations show that this effect is small, although more work should be done to confirm this¹. Finally, there are metal contaminations. For example, Si III that tracks the hydrogen fluctuations contaminates the Ly- α forest flux measurements at separation of 2271 km/s. The cross-correlation between Ly- α forest absorption and Si III absorption, if misinterpreted as Ly- α –Ly- α correlations could potentially bias the BAO measurements [38, 42]. Further contamination arises where the metal absorption traces large scale structures at a significantly different redshift. For example C IV traces structure at $z = 1.7$ at wavelengths which probe the Ly- α forest at $z = 2.4$ [48]. For BAO measurements these can be reliably corrected by including them as a part of the model. For other uses, such as broadband power spectrum measurements, a combination of nuisance modeling, accurate mock spectra and numerical simulations should remove any potential

¹There is an additional effect because Ly- α quasar lines of sight terminate in quasars, which are themselves tracers of the underlying structure, but this can be explicitly shown to be a small effect.

biases associated with these complications.

Perhaps the most important systematic effects will come from imperfections in the instrument and data reduction. For example, artificial features in the mean transmission at the position of galactic Balmer transitions were noticed in BOSS data [39]. These were tracked down to the imperfect interpolation in calibration vectors when these features were masked in calibration stars. Although such effects are on average calibrated out, they can in principle produce sharp features in correlation at certain pairs of wavelengths that could potentially contaminate the BAO measurements. Other effects include noise calibration and its Poisson nature, imperfect sky subtraction, etc. Fortunately, there are no fundamental obstacles to modeling the listed systematics with a carefully executed pipeline. The sheer amount of data that will be available and the relatively high signal-to-noise of true small scale fluctuations in the forest will allow us to check the data in many different ways and validate the data reduction pipeline.

2.3 Measuring Growth of Structure with Redshift Space Distortions

DESI will observe redshifts, which reflect the velocities due to expansion, but also the peculiar velocities due to gravitational attraction by large scale structure. Peculiar velocities are observable in redshift surveys because they alter the correlations between galaxies along the line of sight, resulting in an anisotropy in the observed clustering. Comparing the expansion history and the growth of large scale structure from redshift space distortions will allow DESI to test General Relativity.

2.3.1 Theory

Galaxies and quasars are point tracers of the underlying cosmic structure. The physics of how they trace the dark matter fluctuations is well understood based on arguments about locality of galaxy formation [49, 50, 51]. On very large scales bias is scale independent and redshift-space distortions are described by linear perturbation theory. Beyond-linear perturbative corrections can be used on intermediate scales before perturbation theory breaks down entirely on small scales [52, 53, 54].

The measurement of the growth of structure relies on redshift-space distortions seen in galaxy surveys. Even though we expect the clustering of galaxies in real space to have no preferred direction, galaxy maps produced by estimating distances from redshifts obtained in spectroscopic surveys reveal an anisotropic galaxy distribution. The anisotropies arise because galaxy redshifts, from which distances are inferred, include components from both the Hubble flow and peculiar velocities driven by the clustering of matter. Measurements of the anisotropies allow constraints to be placed on the rate of growth of clustering [55, 56].

On large scales, the observed large-scale structure is basically described by a small fractional perturbation $\delta(\mathbf{x}) = \delta\rho(\mathbf{x})/\bar{\rho} = (\rho(\mathbf{x}) - \bar{\rho})/\bar{\rho}$ to the uniform density. Ignoring the higher-order contributions, the perturbation in redshift space (δ_s) is related to the real space perturbation at directional cosine μ between line-of-sight direction and the wave-number \mathbf{k} , by the Kaiser relation [57],

$$\delta_s(\mathbf{k}) = \delta(\mathbf{k})(1 + \beta\mu^2) \quad (2.9)$$

Here $\beta = f/b$, where b is the galaxy bias and f is related to the linear growth function $D(a)$ by

$$f = \frac{d \ln D(a)}{d \ln a}. \quad (2.10)$$

In the linear regime, density perturbations grow proportional to $D(z)$ which increases with decreasing z .

In GR, $D(z)$ is completely specified by the expansion history even in the presence of dark energy; this is no longer generically true in alternative theories of gravity. The behavior of f in GR is given, to a good approximation, by

$$f \simeq \Omega_m(z)^\gamma, \quad (2.11)$$

where γ is the growth index, approximately 0.55 in GR, and where $\Omega_m(z)$ is the fraction of the total energy density in the form of matter at redshift z . In alternative gravity theories, a common simple parameterization of the modified growth rate is to alter the growth index γ . [58] demonstrated that a DESI-like survey could constrain γ to 0.04 (7%). More general modifications might involve modifying (in a time- and scale-dependent manner) the potentials that enter the metric. Precise growth measurements over a wide range of redshifts and scales, combined with constraints from overlapping CMB and weak lensing surveys, make large galaxy surveys like DESI excellent probes of gravity (see [59] for a recent review). Here, we focus on scale-independent growth rates for large-scale structure, but the DESI data set will allow more complicated investigations.

As an important example of extensions, we highlight the Bright Galaxy Survey, where we will be mapping a smaller volume ($z < 0.4$) at substantially higher number density and with more diversity of galaxies. This redshift range is crucial because it is when dark energy dominates and any associated modifications of gravity would be expected to be strongest. Getting the best precision out of this limited volume requires spectroscopy to produce a 3-D map of the density field. The BGS will test for modifications of gravity directly via the redshift-distortion method, including the novel methods of using multiple tracers in order to suppress sample variance [60]. But the search can be extended via spectroscopic detection of clusters and groups, along with galaxy halo occupation modeling, to measure the amplitude of clustering by halo abundances [61, 62]. The maps can also be correlated with weak lensing maps (e.g., from DES, LSST, Euclid, or CMB-S4) to measure the amplitude of clustering [63, 64]. Comparing the observed velocity field to the expected velocity field sourced from the lensing matter overdensities enables further tests of modified gravity models of cosmic acceleration [55]. Finally, the more detailed map will allow tests of screening theories on smaller scales [65, 66], in which one considers the response of individual galaxies to the predicted gravitational field.

In the Kaiser approximation, the redshift space power spectrum, P_s , is given by

$$P_s(\mathbf{k}) = (b + f\mu^2)^2 P_m(k) \quad (2.12)$$

where P_m is the linear theory mass power spectrum. In principle, this prescribed anisotropy provides a means of measuring f , and through it the growth of gravitational structures. However, in the above, the measurements of f are degenerate with the amplitude of the matter power spectrum. Therefore the combination $f(z)\sigma_8(z)$ is the actual observable, where the normalization of the power spectrum $P(k)$ is proportional to $\sigma_8^2(z)$ ².

2.3.2 Systematics

Galaxies are expected to follow the same gravitational potential as the dark matter and hence have the same velocities. The main theoretical systematic uncertainty in RSD is that nonlinear velocity effects extend to rather large scales and give rise to a scale-dependent and angle-dependent clustering signal. It is easy to see these effects in any real redshift survey: one sees elongated features along the line of sight, called the Fingers of God (FoG). The FoG are caused by random velocities inside virialized objects such as clusters, which scatter galaxies along the radial direction in redshift space, even if they have a localized spatial position in real space. This is just an extreme example and other related effects, such as nonlinear infall streaming motions, also cause nonlinear corrections. In addition, RSD measure velocities as sampled at the galaxy positions. One is thus probing not the velocity field, but rather the momentum density field. Galaxies are a biased tracer of the dark matter and this introduces scale dependent effects into RSD statistics even if galaxies are simply a linear tracer of the dark matter.

There are a plethora of approaches [67, 68, 69, 52, 53, 54] to modeling redshift space distortions in the literature, and the analyses in Table 2.1 make use of many of them. It has been firmly established that the Kaiser formula is inadequate to recover information faithfully on the quasilinear scales of interest, and so most analyses now adopt some form of perturbative corrections. However, because these corrections depend strongly on the halo bias [70, 71], methods calibrated on purely the dark matter power spectrum are of limited utility. Moreover, the details of the mapping between galaxies and dark matter halos also strongly modify the correlation function, mostly through FoG

² σ_8^2 is defined to be the variance of the matter density field averaged in spheres of $8 h^{-1}\text{Mpc}$ and traditionally used to parametrize the amplitude of the power spectrum.

effects. All of these effects can induce 10% effects on RSD at $k \sim 0.1 \, h/\text{Mpc}$. Current models of RSD are able to reproduce these nonlinear effects at the percent level for $k < 0.05\text{--}0.1 \, h/\text{Mpc}$. Extending this to smaller scales would increase the power of the DESI RSD survey. This will require us to improve our bias models and the realism of our simulations.

Most of the observational systematics examined in detail in the SDSS-III BOSS [see 11] primarily affect clustering on the largest scales; currently these are of little concern for RSD measurements, for which the signal comes primarily from the smallest scales included in the measurements. The most important systematic effect is the estimate of a survey’s radial selection function [72, 11]. Since the redshift distribution of targets cannot be predicted precisely a priori, it must be measured directly from the observed galaxies’ redshift distribution. Doing so removes some cosmological radial modes from the observed galaxy overdensity field, resulting in a bias in the monopole-quadrupole amplitudes at the $< 0.2\sigma$ level. The ratio of systematic to statistical uncertainty should remain relatively constant with survey area for a given redshift distribution, since the statistical errors on the correlation function and $n(z)$ shrink at the same rate.

2.3.3 Current Status of RSD Measurements

Redshift-space-distortion measurements have now been performed on a host of surveys, which we summarize in Table 2.1 and show in the left panel of Figure 2.4; taken together, these surveys provide a measure of the growth rate of cosmic structure good to about 3% in the low redshift Universe. Almost all of these measurements of $f\sigma_8$ are derived from the anisotropy in the two-point correlations of the observed galaxy density field. The anisotropic correlation from SDSS-III BOSS DR11 CMASS sample is shown in Figure 2.5. While there have been some analyses directly on the two-dimensional correlation function $\xi(r_p, r_\pi)$ [e.g., 73, 74, 75, 76], most authors further compress the data into multipoles [e.g., 77, 72, 78, 79, 80] or wedges [81, 82]. Efficient information compression is necessary when the covariance matrix of the observables are estimated from a finite number of mock surveys [83].

Most of these measurements assume a flat Λ CDM cosmology to model the redshift-distance relation (see [76] for an exception); dropping this assumption degrades the measurement of $f\sigma_8$. However, the combination of geometric and dynamical constraints available from the analysis of anisotropic galaxy clustering is quite complementary to isotropic BAO measurements for constraining dark energy. For instance, in the case of SDSS-III BOSS DR11 for a flat w CDM cosmology, the combination of Planck and the BOSS BAO measurements constrain $w = -1.01 \pm 0.08$ [6], while including the geometric and dynamical information in the quadrupole correlation function (term proportional to μ^2) yields $w = -0.993 \pm 0.056$ [80].

Considering instead tests of gravity given a “known” expansion history, Figure 2.4 shows that for a flat Λ CDM cosmology in general relativity, the predicted redshift evolution of the observable $f\sigma_8$ is quite mild in the redshift range that has been studied observationally. These observations can begin to distinguish between gravity models ($f(R)$ and DGP are shown), though there is still substantial uncertainty in the theoretical predictions simply due to uncertainties in both the matter density Ω_m and overall matter power spectrum normalization, σ_8 . The right-hand side of Figure 2.4 shows constraints in the Ω_m - γ plane from BOSS DR11 [80]. These data yield a 16% constraint on the growth index. DESI will improve on the precision of the growth constraint from all previous measurements by a factor of $\sim 4\text{--}10$ [84], depending on advances in analysis and theoretical modeling. In addition, it will provide measurements to significantly higher redshifts.

Two surveys in particular are pathfinders for DESI targets: WiggleZ [88] analyzed emission line galaxies with bias b near 1, while SDSS-II and SDSS-III BOSS study luminous red galaxies (LRGs) with a bias near 2. WiggleZ included much smaller scales in their RSD analysis, which led

Table 2.1: Compilation of RSD-based $f\sigma_8$ measurements from [85]. For the BOSS DR11 galaxy sample we cite the measurement of [80]. Other analyses of DR11 find consistent results [82, 79]

z	$f\sigma_8$	survey	reference
0.067	0.42 ± 0.06	6dFGRS	[75]
0.17	0.51 ± 0.06	2dFGRS	[86]
0.22	0.42 ± 0.07	WiggleZ	[77]
0.25	0.35 ± 0.06	SDSS LRG	[72]
0.37	0.46 ± 0.04	SDSS LRG	[72]
0.41	0.45 ± 0.04	WiggleZ	[77]
0.57	0.45 ± 0.03	BOSS CMASS	[80]
0.6	0.43 ± 0.04	WiggleZ	[77]
0.77	0.49 ± 0.18	VVDS	[87]
0.78	0.38 ± 0.04	WiggleZ	[77]

to impressive constraints given the number of galaxies in the survey. However, they were not able to generate easily a large N -body simulation volume capable of resolving the halos expected to host emission line galaxies, and so their theoretical modeling is necessarily less well-tested. By comparison, LRGs are hosted by massive halos that can easily be simulated. The perturbation-based model of [78] was carefully calibrated against N -body-based mock-galaxy catalog and included realistic effects like the “Fingers-of-God” (the elongated structure in the right panel of Figure 2.5). However, because these effects are so strong, their analysis was restricted to relatively large scales.

Ongoing progress in combining the perturbative analytic results with those of N -body simulations should pave way for the increased theoretical prediction accuracy necessary to extract RSD information at small spatial scales




Figure 2.4: *Left:* The data points show the CMASS DR11 measurement of $f\sigma_8$ (gold pentagon; [80]) along with similar, low redshift, measurements and 1σ error bars as presented in Table 2.1. The three stripes show theoretical predictions for different gravity models allowing for uncertainty in the background cosmological parameters, constrained using only the WMAP 7 data [89]. Figure adapted from [85]. *Right:* Joint constraints in the Ω_m - γ plane from BOSS DR11, where γ is the growth index of structure, as defined in Eq. (2.11). Figure taken from [80].




Figure 2.5: The two-dimensional correlation function of the BOSS DR11 CMASS galaxies, measured perpendicular (x-axis) and parallel (y-axis) to the line of sight. The BAO ring, distorted by redshift space distortions is clearly visible, as is the characteristic squashing of the correlation function on large scales.

2.4 Distance, Growth, Dark Energy, and Curvature Constraint Forecasts

DESI’s observational program defined in the Requirements Document and described in this Report specifies the numbers of galaxies and Ly- α forest sources and their distribution that will be measured. Using the specified quality of those observations, we can predict the precision with which cosmological parameters will be determined by DESI. Thanks to the unprecedented scope of DESI’s spectroscopic measurements, these measurements will take us to a new level — Stage-IV — in cosmological exploration.

2.4.1 Forecasting Overview

We use the Fisher matrix formalism to estimate the parameter constraining power of the finished survey, largely following [84]. Our baseline cosmological model is flat Λ CDM. This model is specified by seven parameters, which are listed together with their fiducial values in Table 2.2. Parameter symbols have their conventional meanings. Our standard fiducial parameter values follow *Planck* results, specifically the P+WP+highL+BAO (P from *Planck*, WP from *WMAP*, highL from high resolution CMB experiments like ACT and SPT) column of Table 5 of [13]. In addition to the conventional six parameters of the minimal cosmological model, we also always vary the amount of tensor modes; however this is largely irrelevant because the T/S measurement is completely dominated by *Planck* and essentially uncorrelated with other parameters.

Isolating the BAO feature gives the most robust, but also most pessimistic, view of the information that one can recover from galaxy clustering measurements, since BAO can be measured even in the presence of large unknown systematic effects (very generally, these will not change the BAO scale [19]). We quote errors on the transverse and radial BAO scales as errors on $D_A(z)/s$ and $H(z)s$, respectively, where s is the BAO length scale. For galaxy and quasar clustering, these measurements are correlated at each redshift with a correlation coefficient of 0.4.

We also quote errors on an isotropic dilation factor R/s , defined as the error one would measure on a single parameter that rescales radial and transverse directions by equal amounts. In this case, for a small change in R , the corresponding variations in the model values of D_A and H are

$$D_A = \left(1 + \frac{\delta R}{R_{\text{fid}}}\right) D_{A,\text{fid}} \quad (2.13)$$

and

$$H = \left(1 + \frac{\delta R}{R_{\text{fid}}}\right)^{-1} H_{\text{fid}} \quad (2.14)$$

where $D_{A,\text{fid}}(z)$ and $H_{\text{fid}}(z)$ are the angular diameter distance and Hubble parameter in a fiducial Universe. An explicit definition of R in terms of the measured H and D_A is generally not needed and depends on the experimental scenario. The simplest cases are easy to understand: for a purely transverse measurement (e.g., photometric survey) $R = D_A$, while for a purely radial measurement (e.g., something closer to the Ly- α forest, although it is not purely radial) $R = H^{-1}$ (or $R = H^{-1}H_{\text{fid}}D_{A,\text{fid}}$, if one is concerned about inequivalent units). For intermediate cases like typical galaxy clustering, the appropriate combination of H and D_A can always be determined given the covariance matrix between them. For example, it is approximately proportional to $D_V(z) \equiv ((1+z)D_A)^{2/3}(cz/H(z))^{1/3}$ in analyses of spherically averaged clustering, such as from 6dF, BOSS, and WiggleZ.

Going beyond BAO, we use “broadband” galaxy power, i.e. measurements of the power spectrum as a function of redshift, wavenumber and angle with respect to the line of sight. This treatment automatically recovers all available information from the two-point clustering, i.e. not

Table 2.2: Parameterization of the cosmological model and parameter values for the fiducial model. The seven parameters in the upper part of the table are always free. Parameters in the second half of the table are extensions of the simplest model discussed below.

Parameter	Value	Description
ω_b	0.02214	Physical baryon density $\omega_b = \Omega_b h^2$ $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$
ω_m	0.1414	Physical matter density $\omega_m = \Omega_m h^2$ (including neutrinos which are non-relativistic at $z = 0$)
θ_s	0.59680 degrees	Angular size of sound horizon at the surface of last scattering acting as a proxy for Hubble's constant
A_s	2.198×10^{-9}	Amplitude of the primordial power spectrum at $k = 0.05 \text{ Mpc}^{-1}$ (for the numerical Fisher matrix we actually use $\log_{10} A_s$)
n_s	0.9608	Spectral index of primordial matter fluctuations with $P(k) \propto k^{n_s}$
τ	0.092	Optical depth to the last scattering surface assuming instantaneous reionization.
T/S	0	Ratio of tensor to scalar perturbations (we assume inflationary tensor fluctuation's spectral index $n_t = -\frac{1}{8}T/S$)
w_0	-1	Equation of state of dark-energy $p = w\rho$
w_a	0	Variable equation of state of dark energy of the form $w = w_0 + (1 - a)w_a$
Ω_k	0	Curvature of the homogeneous model
α_s	0	Running of the spectral index $\alpha_s = d \log n_s / d \log k$ with pivot scale $k = 0.05 \text{ Mpc}^{-1}$
Σm_ν	0.06 eV	Sum of neutrino masses (we assume they are degenerate)
$N_{\nu, \text{eff}}$	3.04	Effective number of neutrino species ($N_{\nu, \text{eff}} > 3.04 \rightarrow$ dark radiation).

just the shape of the isotropic power spectrum, but also redshift-space distortions, Alcock-Paczynski [90], and the BAO information.

The broadband Fisher matrix is calculated by combining the inverse variance of the power spectrum $P(\mathbf{k})$ of each Fourier mode with the derivative of power in each mode with respect to set of cosmological parameters. We divide the survey into a set of redshift slices and coadd the resulting matrices. The model for the three-dimensional power spectrum of the galaxy or Ly- α distribution is

$$\tilde{P}(k, \mu, z) = b(z)^2 (1 + \beta(z)\mu^2)^2 P_{\text{mass}}(k, z) D(k, \mu, z), \quad (2.15)$$

where μ is the angle of the wavevector to the line of sight, k is the wavenumber, b is the linear bias parameter, β the redshift space distortion parameter and $D(k, \mu, z)$ is a non-linear correction calibrated from simulations (for the Ly- α forest this is given by [91] and for galaxies it is based on the information damping factors of [92]). The Fisher matrix calculation will integrate over all μ and a suitable range of k . The inverse variance of the power spectrum of each mode gets contributions from both the intrinsic sample variance and the shot noise. This results in an effective volume $V_{\text{eff}}(\tilde{P})$ of each redshift slice that is given by $V_{\text{eff}}(\tilde{P}) = [1 + 1/(n\tilde{P})]^{-2} V_{\text{survey}}$ [93]. The value $n\tilde{P}$ represents the ratio of true clustering power to that from shot noise. Alternatively, it can be seen as the signal-to-noise ratio per mode (redshift, wavenumber, and orientation slice): if $n\tilde{P} > 1$ then

roughly the signal exceeds the sample variance uncertainty for that mode.

For the galaxy survey, we use large-scale broadband power up to some quoted k_{\max} . At small scales, $k > k_{\max}$, we continue to use BAO information. We use two simple choices of k_{\max} : $0.1 \, h \, \text{Mpc}^{-1}$ and $0.2 \, h \, \text{Mpc}^{-1}$. These cutoffs are intended to indicate sensitivity of results to the effective scale where information is recovered after making corrections for non-linearity, after marginalization over suitable non-linear bias parameters. It will be a major program of the next decade to figure out exactly how to do this fitting in practice for a high precision survey like DESI; how well we can do this will determine how well we can measure parameters. As discussed in [84], $k_{\max} \sim 0.1 \, h \, \text{Mpc}^{-1}$ corresponds roughly to the performance of current analyses, while $k_{\max} \sim 0.2 \, h \, \text{Mpc}^{-1}$ is more of a stretch goal for the DESI era (some improvement over current analysis can be expected simply by going to higher redshift where the non-linear scale is smaller).

The redshift-space distortions can effectively constrain two parameter combinations, $b(z)\sigma(z)$ and $f(z)\sigma(z)$, where $\sigma(z) \propto P_{\text{mass}}^{1/2}(z, k)$ is the RMS normalization of the linear mass density fluctuations as a function of z . In Table 2.3, we quote projected constraints on $f\sigma$ for different maximum k assumptions e.g., $f\sigma_{0.1}$ means the error calculation included information up to $k_{\max} = 0.1 \, h \, \text{Mpc}^{-1}$. These fractional errors are equivalent to what one usually sees quoted as an error on “ $f\sigma_8$ ”. The $f\sigma_k$ precision we project for DESI, aggregated over all redshifts, is $\sim 0.74\%$ for $k_{\max} = 0.1 \, h \, \text{Mpc}^{-1}$, or $\sim 0.38\%$ for $k_{\max} = 0.2 \, h \, \text{Mpc}^{-1}$.

2.4.2 Baseline Survey

Our baseline assumption for science projections is that DESI runs over an approximately five-year period covering $14,000 \, \text{deg}^2$ in area. DESI will target four types of objects: Bright Galaxies (BGS), Luminous Red Galaxies (LRGs), Emission Line Galaxies (ELGs) [94], and quasars. Details on how these objects are targeted can be found in Section 3. In what follows, most calculations are done for this baseline survey. We additionally provide several relevant calculations for the required minimum survey with the same target number densities over $9,000$ instead of $14,000 \, \text{deg}^2$ in area.

The number densities used here, plotted in Figure 2.6, are based on the selection criteria for each object type described in the following chapter. We assume fiducial biases follow constant $b(z)D(z)$, where $D(z)$ is the linear growth factor normalized by $D(z=0) \equiv 1$. For LRGs we use $b_{\text{LRG}}(z)D(z) = 1.7$. For ELGs we use $b_{\text{ELG}}(z)D(z) = 0.84$ [94]. For quasars we use $b_{\text{QSO}}(z)D(z) = 1.2$ (loosely based on [95]). For the BGS, we use $b_{\text{BGS}}(z)D(z) = 1.34$, but the results are insensitive to this value because of the much higher number density in most of the BGS volume. Note that these forms keep the observed clustering amplitude of each individual tracer constant with redshift, in agreement with observations (more detailed references for bias evolution are given below, in sections 3.1, 3.2, 3.3.1, and 3.4.1 for BGS, LRGs, ELGs, and QSOs, respectively).

The signal-to-noise for typical BAO-scale modes in redshift space is shown in Figure 2.7, along with the same quantity computed for several other experiments for comparison [84].

We evaluate $\bar{n}P$ at $k = 0.14 \, h \, \text{Mpc}^{-1}$, $\mu = 0.6$, an approximate center-of-weight point for BAO measurements. We chose these values by looking for the point where $\bar{n}P = 1$ corresponded to the optimum in a trade-off between area and number density at fixed total number of objects (specifically, for the full range of parameters covered by DESI LRGs and ELGs). This definition reflects the origin of the idea that $\bar{n}P = 1$ is a special point, but it should be kept in mind that achieving $\bar{n}P$ by this definition does leave a survey significantly farther away from the sample variance limit than the traditional definition $k = 0.2 \, h \, \text{Mpc}^{-1}$, $\mu = 0$.

The spectral signal-to-noise ratio that we use, computed using the `bbspecsim` code [96], is shown in Figure 2.8.




Figure 2.6: DESI number densities, per unit z , per square degree, used in cosmology projections (Table 2.3 and 2.7).




Figure 2.7: Signal to noise comparison of the DESI galaxy survey against other precursor (Stage II and Stage III) and upcoming (Stage IV) spectroscopic surveys. Shown is $\bar{n}P(k = 0.14 h \text{ Mpc}^{-1}, \mu = 0.6)$. The DESI forecasts do not include the Ly- α forest contribution. Including this would give an effective $\bar{n}P \sim 0.3$ at $z \sim 2.5$. Note that the large area covered by DESI provides an advantage reflected in Figure 2.9.




Figure 2.8: Signal-to-noise ratio per Å used for DESI quasar spectra (detector noise, not absorption noise), for different g magnitudes, accounting for mean Ly- α forest absorption.

2.4.3 Summary of Forecasts

Table 2.3 lists the basic galaxy and quasar BAO distance measurement projections, and RSD $f(z)\sigma_8(z)$ error projections for two different k_{\max} values for our baseline 14K survey. We provide the same set of calculations in Table 2.4 for our threshold 9K survey. Tables 2.5 and 2.6 shows the projections for the Bright Galaxy Survey for 14K and 9K square degrees, respectively. Table 2.7 lists the Ly- α forest BAO distance measurement projections, including cross-correlations with quasars in the same redshift range for a $z > 1.9$ Ly- α forest survey; 2.8 presents the same calculations for the threshold 9K survey. The BAO errors are also shown in Figure 2.9, along with those from other experiments for comparison (see [84] for a description of the other experiments).

Table 2.3: Summary of forecasted constraints achievable by DESI, covering 14,000 deg². Indications of signal to noise, nP , are given at two values of $k, \mu = \{0.2 \text{ h Mpc}^{-1}, 0\}$ and $\{1.4 \text{ h Mpc}^{-1}, 0.6\}$. The fractional error on the normalization of $f(z)P^{1/2}(k, z)$ is $\sigma_{f\sigma_k}/f\sigma_k$, assuming known shape of the power spectrum and known geometry, using $k_{\max} = k \text{ h Mpc}^{-1}$. The dilation factor R is defined to be a parameter rescaling the radial and transverse distances by equal factors.

z	$\frac{\sigma_{R/s}}{R/s}$ %	$\frac{\sigma_{DA/s}}{DA/s}$ %	$\frac{\sigma_{Hs}}{Hs}$ %	$\bar{n}P_{0.2,0}$	$\bar{n}P_{0.14,0.6}$	V [h ⁻¹ Gpc ³]	$\frac{dN_{ELG}}{dz \text{ ddeg}^2}$	$\frac{dN_{LRG}}{dz \text{ ddeg}^2}$	$\frac{dN_{QSO}}{dz \text{ ddeg}^2}$	$\frac{\sigma_{f\sigma_{0.1}}}{f\sigma_{0.1}}$ %	$\frac{\sigma_{f\sigma_{0.2}}}{f\sigma_{0.2}}$ %
0.65	0.57	0.82	1.50	2.59	6.23	2.63	309	832	47	3.31	1.57
0.75	0.48	0.69	1.27	3.63	9.25	3.15	2269	986	55	2.10	1.01
0.85	0.47	0.69	1.22	2.33	5.98	3.65	1923	662	61	2.12	1.01
0.95	0.49	0.73	1.22	1.45	3.88	4.10	2094	272	67	2.09	0.99
1.05	0.58	0.89	1.37	0.71	1.95	4.52	1441	51	72	2.23	1.11
1.15	0.60	0.94	1.39	0.58	1.59	4.89	1353	17	76	2.25	1.14
1.25	0.61	0.96	1.39	0.51	1.41	5.22	1337	0	80	2.25	1.16
1.35	0.92	1.50	2.02	0.22	0.61	5.50	523	0	83	2.90	1.73
1.45	0.98	1.59	2.13	0.20	0.53	5.75	466	0	85	3.06	1.87
1.55	1.16	1.90	2.52	0.15	0.40	5.97	329	0	87	3.53	2.27
1.65	1.76	2.88	3.80	0.09	0.22	6.15	126	0	87	5.10	3.61
1.75	2.88	4.64	6.30	0.05	0.12	6.30	0	0	87	8.91	6.81
1.85	2.92	4.71	6.39	0.05	0.12	6.43	0	0	86	9.25	7.07

Table 2.4: Like Table 2.3, except with DESI covering only 9,000 deg².

z	$\frac{\sigma_{R/s}}{R/s}$ %	$\frac{\sigma_{DA/s}}{DA/s}$ %	$\frac{\sigma_{Hs}}{Hs}$ %	$\bar{n}P_{0.2,0}$	$\bar{n}P_{0.14,0.6}$	V h ⁻¹ Gpc ³	$\frac{dN_{ELG}}{dz \text{ ddeg}^2}$	$\frac{dN_{LRG}}{dz \text{ ddeg}^2}$	$\frac{dN_{QSO}}{dz \text{ ddeg}^2}$	$\frac{\sigma_{f\sigma_{0.1}}}{f\sigma_{0.1}}$ %	$\frac{\sigma_{f\sigma_{0.2}}}{f\sigma_{0.2}}$ %
0.65	0.71	1.02	1.87	2.59	6.23	1.69	309	832	47	4.12	1.96
0.75	0.59	0.86	1.58	3.63	9.25	2.03	2269	986	55	2.62	1.26
0.85	0.59	0.86	1.53	2.33	5.98	2.34	1923	662	61	2.64	1.26
0.95	0.61	0.91	1.52	1.45	3.88	2.64	2094	272	67	2.61	1.24
1.05	0.72	1.12	1.70	0.71	1.95	2.90	1441	51	72	2.79	1.39
1.15	0.75	1.17	1.74	0.58	1.59	3.14	1353	17	76	2.80	1.42
1.25	0.76	1.19	1.74	0.51	1.41	3.35	1337	0	80	2.81	1.44
1.35	1.15	1.87	2.52	0.22	0.61	3.54	523	0	83	3.62	2.16
1.45	1.22	1.99	2.66	0.20	0.53	3.70	466	0	85	3.82	2.34
1.55	1.45	2.37	3.14	0.15	0.40	3.84	329	0	87	4.40	2.84
1.65	2.20	3.59	4.74	0.09	0.22	3.95	126	0	87	6.36	4.50
1.75	3.59	5.79	7.86	0.05	0.12	4.05	0	0	87	11.11	8.49
1.85	3.64	5.87	7.97	0.05	0.12	4.13	0	0	86	11.53	8.82

DESI will provide high precision measurements of the Universe's expansion rate over billions of years. Using the Ly- α forest technique, coverage will include the early times when the expansion

Table 2.5: Like Table 2.3, except for the DESI Bright Galaxy Survey, covering 14,000 deg².

z	$\frac{\sigma_{R/s}}{R/s}$ %	$\frac{\sigma_{D_A/s}}{D_A/s}$ %	$\frac{\sigma_{Hs}}{Hs}$ %	$\bar{n}P_{0.2,0}$	$\bar{n}P_{0.14,0.6}$	V [$h^{-1}\text{Gpc}^3$]	$\frac{dN_{BGS}}{dz d\text{deg}^2}$	$\frac{\sigma_{f\sigma_{0.1}}}{f\sigma_{0.1}}$ %	$\frac{\sigma_{f\sigma_{0.2}}}{f\sigma_{0.2}}$ %
0.05	4.33	6.12	12.10	146.60	352.91	0.04	1165	33.24	14.08
0.15	1.66	2.35	4.66	59.47	144.69	0.23	3074	12.47	5.25
0.25	1.07	1.51	2.97	14.84	36.43	0.58	1909	7.69	3.25
0.35	0.91	1.32	2.44	3.21	7.94	1.04	732	5.83	2.60
0.45	1.56	2.39	3.69	0.35	0.87	1.55	120	6.35	3.77

Table 2.6: Like Table 2.5, but for a 9,000 deg² Bright Galaxy Survey.

z	$\frac{\sigma_{R/s}}{R/s}$ %	$\frac{\sigma_{D_A/s}}{D_A/s}$ %	$\frac{\sigma_{Hs}}{Hs}$ %	$\bar{n}P_{0.2,0}$	$\bar{n}P_{0.14,0.6}$	V [$h^{-1}\text{Gpc}^3$]	$\frac{dN_{BGS}}{dz d\text{deg}^2}$	$\frac{\sigma_{f\sigma_{0.1}}}{f\sigma_{0.1}}$ %	$\frac{\sigma_{f\sigma_{0.2}}}{f\sigma_{0.2}}$ %
0.05	5.39	7.63	15.09	146.60	352.91	0.02	1165	41.46	17.56
0.15	2.07	2.93	5.81	59.47	144.69	0.15	3074	15.55	6.54
0.25	1.33	1.89	3.70	14.84	36.43	0.38	1909	9.59	4.05
0.35	1.14	1.64	3.04	3.21	7.94	0.67	732	7.27	3.24
0.45	1.94	2.98	4.60	0.35	0.87	1.00	120	7.92	4.71

rate was decreasing (when the matter density, not the dark energy density, was controlling the rate). In Figure 2.10 we show how DESI will improve these measurements over those existing today.

Table 2.9 shows Dark Energy Task Force (DETF) Figures of Merit (FoMs) [4]. For the common normalization convention that we follow, the FoM is simply $(\sigma_{w_p}\sigma_{w'})^{-1}$ where $w(z) = w_p + (a_p - a)w'$ and a_p is chosen to make the errors on w_p and w' independent. Because the DETF FoM model is defined to include the possibility of curvature, we include curvature projections in Table 2.9. The figure of merit results are reflected in Figure 2.11.

Importantly, Table 2.9 shows that these surveys exceed the Stage IV FoM threshold. We take this to be a value of 110, based on a 10-fold improvement of the value of 11 from [102]. This is the same Stage IV definition that LSST used in their Conceptual Design Report. The 9,000 square degrees DESI survey achieves 121 with galaxies and Ly- α forest BAO. We note that these computations include only BAO and CMB, without even the Stage II Supernovae Ia results from [102]. Including DESI galaxy broadband clustering or other dark energy probes boost the Figure of Merit well above 110.

As this 9000 square degree survey forecast meets the Stage IV threshold and hence the Mission Need, we have adopted it as the quantitative basis for the Level 1 Science Requirement for the DESI project. We aggregate the BAO performance into three redshift ranges, R in $0.0 < z < 1.1$ and $1.1 < z < 1.9$ and H in $1.9 < z < 3.7$, for the L1 requirements, so as to leave flexibility in the exact redshift distribution of targets. An extensive discussion of how the FOM depends on variation in survey parameters was presented in the DESI Conceptual Design Review.

The measurements of $f\sigma_8$ from redshift-space distortion provide the means for testing General Relativity. Figure 2.12 shows the rate of growth of structure, f , as a function of the redshift. Forecasted DESI errors, assuming information at $k < 0.2 h \text{ Mpc}^{-1}$, are shown on the ΛCDM curve. Alternative gravity models generically predict scale-dependent growth, and here we show theoretical expectations for the $f(R)$ modified theory of gravity evaluated at two scales (two values of k), as well as predictions for the DGP braneworld theory. DESI can clearly distinguish between these models.

Table 2.7: $z > 1.9$ Ly- α forest quasar survey, over 14000 sq. deg. Parameter errors are in percent relative to the BAO scale, s .

z	$\frac{\sigma_{R/s}}{R/s} (\%)$	$\frac{\sigma_{DA/s}}{DA/s} (\%)$	$\frac{\sigma_{Hs}}{Hs} (\%)$	$\frac{dN_{QSO}}{dz d\text{deg}^2}$
1.96	1.43	2.69	2.74	82
2.12	1.02	1.95	1.99	69
2.28	1.09	2.18	2.11	53
2.43	1.20	2.46	2.26	43
2.59	1.34	2.86	2.47	37
2.75	1.53	3.40	2.76	31
2.91	1.81	4.21	3.18	26
3.07	2.16	5.29	3.70	21
3.23	2.75	7.10	4.57	16
3.39	3.86	10.46	6.19	13
3.55	5.72	15.91	8.89	9
3.70	-	-	-	7
3.86	-	-	-	5
4.02	-	-	-	3

Table 2.8: Like Table 2.7, except with DESI covering only 9,000 deg².

z	$\frac{\sigma_{R/s}}{R/s} (\%)$	$\frac{\sigma_{DA/s}}{DA/s} (\%)$	$\frac{\sigma_{Hs}}{Hs} (\%)$	$\frac{dN_{QSO}}{dz d\text{deg}^2}$
1.96	1.78	3.35	3.42	82
2.12	1.27	2.43	2.48	69
2.28	1.37	2.72	2.63	53
2.43	1.49	3.07	2.82	43
2.59	1.67	3.57	3.08	37
2.75	1.91	4.24	3.44	31
2.91	2.25	5.26	3.96	26
3.07	2.69	6.60	4.62	21
3.23	3.43	8.86	5.70	16
3.39	4.81	13.05	7.72	13
3.55	7.14	19.85	11.09	9
3.70	-	-	-	7
3.86	-	-	-	5
4.02	-	-	-	3




Figure 2.9: The fractional error on the dilation factor, R , as a function of redshift presented in comparable bins for DESI, BOSS, *Euclid*, *WFIRST*, HETDEX, and eBOSS. This gives an indicative error on distance measurements to each redshift. The forecasts for a 14,000 deg^2 DESI Bright Galaxy Survey (BGS) are also shown. DESI will provide the best measurements over much of the region and is competitive with space-based missions, which will come later. We use 50 million total galaxies for *Euclid*, following their Definition Study Report [97], although recently it has been suggested that this may be optimistic [98].

Table 2.9: DETF Figures of Merit and uncertainties σ_{w_p} and σ_{Ω_k} . σ_{w_p} is the error on w at the pivot redshift, which also equal to the error on a constant w holding $w_a = 0$. σ_{Ω_k} is the error on the curvature of the Universe, Ω_k . All DESI lines contain the BGS, and BOSS in the range $0.45 < z < 0.6$ that does not substantially overlap with DESI. All cases include *Planck* CMB constraints. The pivot point, where $w(a)$ has minimal uncertainty is indicated by a_p . We note that a FoM of 110 is 10 times the Stage II level of [102], which we take to be the definition of Stage IV. DESI BAO galaxy exceeds this threshold even with a 9,000 square degree survey.

Surveys	FoM	a_p	σ_{w_p}	σ_{Ω_k}
BOSS BAO	37	0.65	0.055	0.0026
DESI 14k galaxy BAO	133	0.69	0.023	0.0013
DESI 14k galaxy and Ly- α forest BAO	169	0.71	0.022	0.0011
DESI 14k BAO + gal. broadband to $k < 0.1 \, h \, \text{Mpc}^{-1}$	332	0.74	0.015	0.0009
DESI 14k BAO + gal. broadband to $k < 0.2 \, h \, \text{Mpc}^{-1}$	704	0.73	0.011	0.0007
DESI 9k galaxy BAO	95	0.69	0.027	0.0015
DESI 9k galaxy and Ly- α forest BAO	121	0.71	0.026	0.0012
DESI 9k BAO + gal. broadband to $k < 0.1 \, h \, \text{Mpc}^{-1}$	229	0.73	0.018	0.0011
DESI 9k BAO + gal. broadband to $k < 0.2 \, h \, \text{Mpc}^{-1}$	502	0.73	0.013	0.0009




Figure 2.10: Expansion rate of the Universe as a function of redshift. In the upper plot, the filled blue circle is the H_0 measurement of [99], the solid black square shows the SDSS BAO measurement of [100], the red square shows the BOSS galaxy BAO measurement of [6], the red circle shows the BOSS Ly- α forest BAO measurement of [42], and the red x shows the BOSS Ly- α forest BAO-quasar cross-correlation measurement of [101]. The lower plot shows projected DESI points.




Figure 2.11: The $w_0 - w_a$ plane showing projected limits (68%) from DESI using just BAO and using the broadband (BB) power spectrum. Also shown is the limit from BOSS BAO. *Planck* priors are included in all cases, and DESI includes the BGS and non-redundant part of BOSS. The figure of merit of the surveys is inversely proportional to the areas of the error ellipses.




Figure 2.12: Growth of structure, f , as a function of redshift, showing projected DESI measurements and their ability to discriminate against alternative gravity models, $f(R)$ (whose scale-dependent growth we show evaluated at two different scales) and DGP. The brown (light) error bars at $z < 0.5$ correspond to DESI Bright Galaxy Survey; these are expected to improve when information from the multiple tracers in the BGS is included. Adopted from the Snowmass report on the growth of cosmic structure [59].

2.4.4 Forecasting Details

2.4.4.1 Galaxy and Quasar Clustering

Our treatment of isolated galaxy BAO follows [92], assuming 50% reconstruction, i.e., reduction of the BAO damping scale of [92] by a factor 0.5, except at very low number density, where we degrade reconstruction based on [103].

Bias uncertainty is modeled by a free parameter in each redshift bin, generally of width $\Delta z = 0.1$, for each type of galaxy. Our results are not sensitive to the redshift bin width [84]. For the broadband signal, we use the same information damping factors from [92] as we use for BAO. This is well-motivated from a theoretical point of view as the non-linear clustering suppresses all linear theory information, not just BAO [16]. We also include the reconstruction factor (50% reduction in damping length), assuming that reconstruction will recover non-BAO information as well. See [84] for more discussion.

2.4.4.2 Ly- α Forest

DESI will also probe large-scale structure using the Ly- α forest [104, 38], i.e., the Ly- α absorption by neutral gas in the intergalactic medium in the spectra of high redshift quasars (it may be possible to do even better at faint magnitudes using Lyman-break galaxies [36]). The distribution of intergalactic gas can be used as a complementary tracer to galaxies of the underlying matter distribution for BAO and broadband power spectrum characteristics.

The constraints from the Ly- α forest are difficult to predict accurately, because they require careful simulation [105, 106]. The forecasts described below we believe are a conservative assessment. We limit the application of Ly- β forest data to BAO only (see below), and do not include cross-correlations with quasar density, nor statistics beyond the power spectrum, such as the bispectrum, which are known to be powerful for breaking IGM model degeneracies (e.g., [107]). Finally, we only use the redshift range $z = 2 - 2.7$.

We model the three dimensional power spectrum of Ly- α using Eq. (2.15) and, except as otherwise noted, we use the method of [36] to estimate the errors obtainable by DESI. We use Table I of [32] to model the dependence of b , β , and fitting parameters of D . While these are primarily valid near $z \approx 2.25$, for BAO the model dependence is not significant. For broadband spectra constraints the bias and damping parameters depend on the amplitude and slope of the linear power spectrum, temperature-density relation [108], and mean level of absorption [109], all of which are varied in our Fisher matrix calculations. To help constrain these parameters, we include the one-dimensional power spectrum, which could be measured from the hundreds of existing high resolution spectra [109, 110].

While past projections used the rest wavelength range $1041 < \lambda < 1185 \text{ \AA}$ (following [104]), for the BAO constraints only, we expand the range to include the Ly- β forest and move slightly closer to the quasar, $985 < \lambda < 1200 \text{ \AA}$, reflecting our increasing confidence that we understand the relevant issues well enough to measure BAO across this range [111]. (The Ly- β forest is the wavelength range $\sim 973 - 1026 \text{ \AA}$ where there is Ly- β absorption on top of the Ly- α absorption. This Ly- β absorption corresponds to the same gas we see in the standard Ly- α forest and should provide some extra information, but we simply assume it can be mostly removed as a source of noise and the underlying Ly- α used to measure BAO to shorter wavelengths in each quasar spectrum.) Gains from this enhancement of effective number density (and cross-correlations with quasars) are substantial because the measurement is quite sparse, i.e., in what for galaxies we would call the shot-noise limited regime.

The cross-correlation of quasars with the Ly- α forest [112] provides a complementary measurement of BAO at high redshift. We combine the two probes of structure in the same volume as described in [84]. The correlation of Ly- α absorption in quasar spectra can also provide other cosmological information, beyond BAO: cosmological parameter constraints from the line of sight power spectrum [104, 113, 114, 115], and from the full shape of the three-dimensional clustering [32]. In the projections below we distinguish between Ly- α forest BAO measurements and broadband measurements that include the one-dimensional power spectrum measurement.

2.5 Cosmology Beyond Dark Energy

While the fundamental goal of DESI is the measurement of the expansion rate of the Universe through BAO and RSD, the enormous spectroscopic survey will measure the two-point correlation function and power-spectrum over a broad range of scales and redshifts. These data will open up broader investigations into cosmology and particle physics.

The broadband power spectrum will provide tests of inflation through its scale dependence. Inflation can also be tested through the scale dependence of the bias of dark matter halos, which constrains the primordial non-Gaussianity. The power spectrum will also reflect the damping of structure by free-streaming neutrinos and thereby give a measure of the sum of the neutrino masses, and possibly reveal previously unknown nearly massless species.

2.5.1 Inflation

The inflationary paradigm is the leading explanation for the origin of the fluctuations of primordial density, which in turn seeded the large-scale structure we observe today. In its simplest formulation, inflation predicts perturbations in the initial distribution that are very nearly scale-independent and Gaussian-distributed about the mean. Inflation has been tested primarily with the CMB observations — starting with *COBE* measurements on large scales in the early 1990s and continuing with the increasingly precise *WMAP* and *Planck* measurements in this millennium. However the CMB temperature measurements are not expected to improve greatly after *Planck* (though CMB polarization has a lot to offer, in particular in testing for signatures of inflationary gravity waves). Large-scale structure measurements have become increasingly precise thanks to 2dF, SDSS, and WiggleZ. These complement the CMB measurements in temporal and spatial scales. The next frontier for tests of inflation is large-scale structure. DESI’s unparalleled three-dimensional picture of the evolution of structure will contribute powerfully.

2.5.1.1 Spectral Index and Its Running

Inflation predicts that the primordial spectrum of density fluctuations is nearly a power law in wavenumber k . The power law is specified by the spectral index defined as

$$n_s(k_0) = \left. \frac{d \ln P}{d \ln k} \right|_{k_0} \quad (2.16)$$

where k_0 is some reference scale, typically chosen to be $k_0 = 0.05 \text{ Mpc}^{-1}$. A perfect power law would correspond to a constant n_s ; in reality, inflation also predicts a small “running” with wavenumber parameterized with the parameter $\alpha = dn_s/d \ln k$, again defined at k_0 . The primordial power spectrum can therefore be written as [116]

$$P(k) = P(k_0)(k/k_0)^{n_s(k_0) + \frac{1}{2}\alpha \ln(k/k_0)}. \quad (2.17)$$

The exact Harrison-Zel’dovich primordial spectrum has $n_s = 1$, while inflation predicts slight deviations from unity. Ruling out $n_s = 1$ at a significant level of confidence would strengthen the case for inflation [117]. Recent *Planck* data currently favor $n_s < 1$ at 5σ ; $n_s = 0.968 \pm 0.006$ [118]. The current limit on running of the spectral index obtained by *Planck* is $dn_s/d \ln k = -0.003 \pm 0.007$ (95% CL). Because it is in the regime of linearity for a wide range of k , the Ly- α forest is an excellent complementary probe of α_s .

In Table. 2.10 we present forecasts on inflationary observables obtained with the Fisher-matrix formalism described in Section 2.4.1, applied to the power spectrum obtained from DESI galaxies,

Table 2.10: Projected constraints on inflationary observables obtained by DESI. In all cases, we include constraints from the *Planck* satellite and BAO information from DESI galaxies, quasars and the Ly- α forest. We show the result of including information from the broadband galaxy power spectrum (“Gal”) out to $k_{\text{max}} = 0.1$ and $0.2 \, h \text{Mpc}^{-1}$, and from the Ly- α forest. The numbers in parentheses show the relative improvement over *Planck*. Broadband Ly- α forest constraints include ~ 100 existing high resolution spectra to constrain the IGM model. n_s constraints assume fixed α_s . Both constraints are marginalized over Σm_ν , and the fiducial values are $n_s = 0.963$, $\alpha_s = 0$.

Data	σ_{n_s}	σ_{α_s}
Gal ($k_{\text{max}} = 0.1 h \text{Mpc}^{-1}$)	0.0025 (1.3)	0.005 (1)
Gal ($k_{\text{max}} = 0.2 h \text{Mpc}^{-1}$)	0.0022 (1.5)	0.004 (1.3)
Ly- α forest	0.0029 (1.1)	0.0027 (1.9)
Ly- α forest + Gal ($k_{\text{max}} = 0.2$)	0.0019 (1.7)	0.0019 (2.7)

quasars, and Ly- α forest, combined with CMB data from the *Planck* satellite. The table shows strong constraints on n_s , and improvements up to a factor of three over *Planck* alone, under the assumption that there is no significant running in the spectral index. Achieving these constraints will require excellent control of broad-band systematics in the Ly- α forest and galaxy analyses. But the effort is worthwhile, as these measurements can have far-reaching implications on our understanding of the very early Universe, as we now describe.

For the spectral index, the increased accuracy implies much better constraints on models of inflation. With the DESI+*Planck* constraints, excellent constraints on the spectral index will effectively reduce the allowed region in the plane of n_s and r , the ratio of tensor to scalar modes, to a vertical line pinned at the measured value of n_s . Combining these results with better measurements of the r from the small-scale CMB experiments will lead to much better constraints on inflationary models. Even without the accompanying r measurements, better determination of the spectral index is important: for example, for inflationary potentials $V(\phi) \propto \phi^m$, where ϕ is the inflaton field, the spectral index and the total number of e-folds of inflation N are related via $1 - n_s = (m+2)/(2N)$ [119]. Hence, for this class of models the duration of the inflationary phase would be determined by DESI very precisely.

Implications of the precise measurements of the running of the spectral index α_s are even more impressive. In standard single-field slow-rolling inflationary models, the running of the spectral index is of the order $O((1 - n_s)^2) \sim 1 \times 10^{-3}$ if $n_s \sim 0.96$. This means that DESI will start to approach the region of expected detection in minimal inflationary models. More importantly, a detection of running *larger* than the slow-roll prediction would imply either that inflation involves multiple fields, or a breakdown of the slow roll approximation [120], or else that a non-canonical kinetic term is controlling inflationary dynamics [121]. *Any* detection of the running of the spectral index would represent a significant advance in our understanding of the physics of inflation.

2.5.1.2 Primordial non-Gaussianity

One of the fundamental predictions of the simplest inflationary models is that the density fluctuations in the early Universe that seeded large-scale structure were nearly Gaussian distributed. A single field slow-roll inflation with canonical kinetic energy and adiabatic vacuum predicts very small amount of non-Gaussianity. A violation of any of these conditions, however, may lead to large non-Gaussianity. A simple, frequently studied model is that of non-Gaussianity of the local type, $\Phi = \phi_G + f_{\text{NL}}(\phi_G^2 - \langle \phi_G^2 \rangle)$, where Φ is the primordial curvature fluctuation and ϕ_G is a Gaussian random field. A detection of nonzero f_{NL} would rule out the simplest model of inflation, while a

non-detection at a level of $f_{\text{NL}} < O(1)$ would rule out many of its alternatives.

The tightest existing upper limits on non-Gaussianity have been obtained from observations of the cosmic microwave background by the *Planck* experiment [122]. Recently, a number of inflationary models have been proposed which predict a potentially observable level of non-Gaussianity, these include those from fast-roll inflation [123, 124, 125, 126, 127], quasi-single field inflation [128, 129], warm inflation [130, 131], and non-Bunch-Davies or excited initial states [123, 132, 133, 134]. There are also hybrids of multi-field and non-slow-roll models [135, 136, 137], and the inclusion of isocurvature modes in the non-Gaussian correlations [138, 139, 140]. Improved limits on non-Gaussianity would rule out some of these models. Conversely, a robust detection of primordial non-Gaussianity would dramatically overturn the simplest model of inflationary cosmology, and provide information that would help us significantly improve our understanding of the nature of physical processes in the early Universe.

Until recently, the most powerful methods to place limits on f_{NL} were based on the bispectrum of the CMB. The constraints from CMB data have improved starting from $\sigma(f_{\text{NL}}) \simeq 3000$ with *COBE* [141] to $\sigma(f_{\text{NL}}) \simeq 20$ with *WMAP* [142], to the tight constraint of $\sigma(f_{\text{NL}}) \simeq 5.8$ with *Planck*'s first year data [143] and finally to $\sigma(f_{\text{NL}}) \simeq 5.0$ with the 2015 data from *Planck* [122]. It is therefore impressive and maybe even surprising that a powerful LSS survey such as DESI can provide comparable but highly complementary constraints to *Planck*. Moreover, as we now show, DESI and *Planck* in combination can provide very tight constraints on distinct *classes* of physically motivated inflationary models.

Powerful constraints on non-Gaussianity can come from the effect that it has on the clustering of dense regions on very large scales [144]. Essentially, the bias of dark matter halos assumes a unique, scale-dependent form at large spatial scales in the presence of primordial non-Gaussianity of local type

$$b(k) \equiv b_0 + \Delta b(k) = b_0 + f_{\text{NL}}(b_0 - 1)\delta_c \frac{3\Omega_M H_0^2}{a g(a) T(k) c^2 k^2}, \quad (2.18)$$

where b_0 is the usual Gaussian bias (on large scales, where it is constant), f_{NL} is the parameter that indicates departures from Gaussianity (when $f_{\text{NL}} \neq 0$), $\delta_c \approx 1.686$ is the collapse threshold, $T(k)$ is the transfer function and $g(a)$ is the growth suppression factor. Notice the unique k^{-2} scale dependence in the presence of primordial non-Gaussianity. Since the bias $b(k)$ is readily measured from the correlation function of galaxies or quasars, classes of inflationary models can be tightly constrained. A first application of this method has been presented using the large-scale clustering of quasar and luminous red galaxies (LRG) galaxy data from the Sloan Digital Sky Survey (SDSS) [145]. The result, a non-detection with one sigma error $\sigma(f_{\text{NL}}) \simeq 25$, was (at the time) comparable to the CMB constraints from *WMAP*. DESI will provide constraints competitive, and very complementary, to those from *Planck*, provided that we have systematics under control [146, 147, 148]

Forecasts for DESI indicate that the 1σ error on the local model from DESI alone will be $\sigma(f_{\text{NL}}) \simeq 5$, and about a factor of two better when combined with the final *Planck* temperature and polarization data. From the fundamental physics point of view, these constraints are very exciting, as they probe not only primordial non-Gaussianity but are likely to detect the additional non-Gaussian signal due to late-time nonlinear interactions of the photon-baryon fluid with gravity (with $f_{\text{NL}}^{\text{late}} \simeq \text{few}$ [149, 150]), and thus provide an additional test of cosmology.

More generally, inflationary models predict a range of possibilities for the scaling of the bias $\Delta b \propto k^{-m}$. For example, $m = 2$ for the local model parameterized by f_{NL} as in Eq. (2.18); multi-field inflationary models generically produce $0 < m \lesssim 2$, and models with modifications to the initial quantum state can produce an even stronger scaling with $m = 3$. Because many of these




Figure 2.13: Constraints on the models of primordial non-Gaussianity with “running”, where the usual parameter f_{NL} is promoted to a power-law function of wavenumber, $f_{\text{NL}}(k) = f_{\text{NL}}^*(k/k_*)^{n_{f_{\text{NL}}}}$. The larger contours show constraints on f_{NL}^* and $n_{f_{\text{NL}}}$ from a first analysis that was applied to WMAP 7 data [151]. The size of the red dot shows the 68% C.L. forecast on the joint constraint expected from the combination of the DESI and full *Planck* data sets, based on projections in Ref. [152].

models therefore leave a strong imprint in the clustering of galaxies and quasars, DESI will be able to strongly constrain whole classes of inflationary models. We show an illustration in Figure 2.13, where we present constraints on the models with “running” of non-Gaussianity, where the usual parameter f_{NL} now runs with wavenumber, $f_{\text{NL}}(k) = f_{\text{NL}}^*(k/k_*)^{n_{f_{\text{NL}}}}$. The larger contours show constraints on f_{NL}^* and $n_{f_{\text{NL}}}$ from a first analysis that was applied to WMAP 7 data [151], while the small, red contour shows the 68% C.L. forecast on the joint constraint expected from the combination of the DESI and full *Planck* data sets, based on projections in Ref. [152]. The latter constraint will shrink the area in the $f_{\text{NL}}^* - n_{f_{\text{NL}}}$ plane by about a factor of 100.

To achieve such excellent constraints, the galaxies measured in DESI must have sufficiently large bias, since only for biased tracers is the non-Gaussian scale-dependent clustering revealed. One way to further improve the errors is by combining two tracers of LSS, one with a high bias and one with a low bias. In this case it may be possible to cancel sampling variance, which is the dominant source of error on large scales [153, 154], but due to low number density this will have to include an additional tracer of structure, potentially combining with the LSST and DES data.

More detailed studies of halo mass distribution of BOSS galaxies, combined with numerical simulations of non-Gaussian models [155] as well as studies of how to mitigate the large-angle systematic errors [156, 157, 148] are needed to provide a better definition of the ultimate reach of DESI for non-Gaussianity studies. However it seems certain that DESI constraints will be at least comparable to the best limits from CMB and that they will provide an excellent temporal and spatial complement to the latter.

Table 2.11: Constraints on the sum of neutrino masses from DESI forecasts in combination with constraints from the *Planck* satellite. The experiment combinations are identified as described in the caption of Table 2.10. The last four cases include the information from *Planck* and DESI BAO measurements. Fiducial values are $\Sigma m_\nu = 0.06$ eV, $N_{\nu,\text{eff}} = 3.04$. Σm_ν constraints assume fixed N_ν , while N_ν is marginalized over Σm_ν .

Data	$\sigma_{\Sigma m_\nu}$ [eV]	$\sigma_{N_{\nu,\text{eff}}}$
Planck	0.56	0.19
Planck + BAO	0.087	0.18
Gal ($k_{\text{max}} = 0.1h \text{ Mpc}^{-1}$)	0.030	0.13
Gal ($k_{\text{max}} = 0.2h \text{ Mpc}^{-1}$)	0.021	0.083
Ly- α forest	0.041	0.11
Ly- α forest + Gal ($k_{\text{max}} = 0.2$)	0.020	0.062

2.5.2 Neutrinos

The effects of neutrinos in cosmology are well understood (for a review, see [158]). They decouple from the cosmic plasma when the temperature of the Universe is about 1 MeV, just before electron-positron annihilation. While ultra-relativistic, they behave as extra radiation (albeit not electromagnetically coupled) with a temperature equal to $(4/11)^{1/3}$ of the temperature of the cosmic microwave background. As the Universe expands and cools, they become non-relativistic and ultimately behave as additional dark matter.

2.5.2.1 Neutrino Mass

The mass of neutrinos has two important effects in the Universe [158]. First, as the neutrinos become non-relativistic after the time of CMB decoupling they contribute to the background evolution in the same way as baryons or dark matter, instead of becoming completely negligible as they would if massless (like photons). This affects anything sensitive to the background expansion rate, e.g., BAO distance measurements. Second, the process of neutrinos becoming non-relativistic imprints a characteristic scale in the power spectra of fluctuations. This is termed the ‘free-streaming scale’ and is roughly equal to the distance a typical neutrino has traveled while it is relativistic. Fluctuations on smaller scales are suppressed by a non-negligible amount, of the order of a few percent. This allows us to put limits on the neutrino masses.

From neutrino mixing experiments we know the differences of the squares of masses of the neutrino mass eigenstates. The splitting between the two states with similar masses is $\Delta m_{21}^2 = (7.50 \pm 0.20) \times 10^{-5} \text{ eV}^2$, while the splitting between the highest and lowest masses squared is $\Delta m_{32}^2 = 2.32_{-0.08}^{+0.12} \times 10^{-3} \text{ eV}^2$. Two things are not known: the absolute mass scale, and whether the two states close together are more or less massive than the third state. In what is called the normal hierarchy, the close states are less massive. In this configuration, the lowest possible masses in eV are 0, 0.009, and 0.048, so the minimal sum of neutrino masses is 0.057 eV. In the inverted hierarchy, the minimal masses are 0, 0.048, and 0.049 eV, for a total of 0.097 eV. This is shown in Figure 2.14.

Table 2.11 shows our projected Σm_ν constraints, obtained through Fisher matrix calculations as discussed above and in [84].

With a projected resolution of 0.020 eV, DESI will make a precision measurement of the sum of the neutrino masses independent of the hierarchy and therefore determine the absolute mass scale for neutrinos, a measurement that is otherwise very challenging. Furthermore, if the masses were




Figure 2.14: The two possible neutrino mass hierarchies. Also shown is what fraction of each mass eigenstate corresponds to a neutrino flavor eigenstate. DESI will be sensitive to the sum of the neutrino masses and possibly to the mass hierarchy.

minimal and the hierarchy normal, DESI would be able to exclude the inverted hierarchy at 2σ .

2.5.2.2 Dark Radiation (e.g., sterile neutrinos)

The other parameter relevant for neutrino physics is the effective number of neutrino species $N_{\nu,\text{eff}}$, which parameterizes the energy density attributed to any non-electromagnetically interacting ultrarelativistic species (including e.g. axions) in units of the equivalent of one neutrino species that fully decouples before electron-positron annihilation. Extra radiation shifts the redshift of matter radiation equality and changes the expansion rate during the CMB epoch, although it does not significantly affect the Universe at the epoch probed by DESI. The value for the standard cosmological model is $N_{\nu,\text{eff}} = 3.04^3$ [159]. The detection of any discrepancy from the expected value would be a truly major result, as it would indicate a sterile neutrino [160], a decaying particle [161], a nonstandard thermal history [162], or perhaps that dark energy does not fade away to $\sim 10^{-9}$ at the time of recombination as expected for the cosmological-constant model [163]. All of these possibilities represent important extensions of the standard cosmological model, and uncovering them would present a major advance of our understanding of the Universe. Our forecasts for this parameter are also shown in Table 2.11. Again we see that the effective number of neutrino species will be measured to $\sim 10\%$ or better, providing strong constraints on the alternative models involving extra sterile neutrinos, axions or partly thermalized species.

In Figure 2.15 we show the improvement in the measurement of several fundamental parameters from cosmology and neutrino physics. The standard is taken to be the results from BOSS together with *Planck*. Displayed is the ratio of the uncertainty from BOSS over the uncertainty from DESI, with *Planck* always included.

³The small increase with respect to $N_\nu = 3$ is due to the fact that some neutrinos are still coupled at the onset of electron-positron annihilation.




Figure 2.15: Improvement in the measurements of w_p , $w' = w_a$, Ω_k , Σm_ν the sum of the neutrino masses, n_s the spectral index, α_s the running of the spectral index, and $N_{\nu,\text{eff}}$ the number of neutrino-like (relativistic) species.

2.6 The Milky Way Survey: Near-Field Cosmology from Stellar Spectroscopy

During conditions unusable for faint galaxy work, DESI will pursue the Bright Galaxy Survey, mapping 10 million galaxies to $z \sim 0.4$ in pursuit of the clustering analyses, such as from BAO and RSD, as described earlier in this chapter. As detailed in section 3.1, the areal density of these bright galaxies is comparable to the fiber density of DESI. Achieving a high completeness in the face of clustering and Poisson fluctuations requires multiple visits, leading to an excess of fibers compared to targets. Indeed, some fibers will be unable to reach a viable galaxy target even on the first pass, and this fraction increases on subsequent passes.

Bright stars are the natural secondary target, and we expect that any bright galaxy survey with the DESI fiber positioner will produce a very large sample of stars as a by-product. This sample is also of high science interest, leading to the definition of the Milky Way Survey. At 17th magnitude, even a short (8-10 min) DESI exposure measures an excellent spectrum with $S/N = 25$ per pixel, which will yield the radial velocity to a few km/s precision and the metallicity. We expect the BGS to generate at least 10 million such spectra. Spectroscopy of individual stars provides radial velocity, effective temperature, surface gravity, chemical abundance distribution, and approximate age. The assembly history of the Milky Way is encoded in the spatial distributions, kinematics, and chemical composition of the various distinct Galactic stellar populations. This information can test cosmological predictions for how galaxies like the Milky Way form and evolve on small scales that are difficult or impossible to test elsewhere in the Universe, and provide a critical test of the small-scale predictions of the Λ CDM model.

The European Space Agency *GAIA* satellite has been successfully launched and will provide a catalog of parallaxes, proper motions, and spectrophotometry for a billion point sources down to $V \sim 20$ over the whole sky. The satellite's RVS spectrograph will supplement those data with radial velocities for millions of brighter stars, although the flux limit is still under investigation due to higher than expected scattered light. DESI can substantially enhance the science return from *GAIA* by providing radial velocities and metallicities for stars much fainter than what the *GAIA* spectrograph can provide. While other projects are planned for spectroscopic follow-up of *GAIA* stars, DESI's higher multiplex, wide field of view, and extremely rapid reconfiguration give it a clear advantage.

The stellar program will put exceptional new constraints on the distribution of dark matter in the Milky Way, a vital measurement that links Galactic science, galaxy formation and cosmology. The Milky Way gravitational potential can be probed via the rotation of the Milky Way beyond 15 kiloparsecs, the motions of newly discovered tidal streams, and the kinematics of bright stars in the distant stellar halo. The uncertainty in the Milky Way mass, density profile, and internal structure currently are critically important systematics in the interpretation of direct and indirect dark matter searches, and the measurements possible with the stellar program will substantially reduce these uncertainties.

Joint metallicity and velocity distribution functions for stars far beyond the solar neighborhood will reveal the recent assembly history of the outer disk and vastly improve our understanding of the structure and formation of the thick disk. The first-ever deep spectroscopic survey of halo main-sequence turn-off stars to 30 kiloparsecs can be used to reconstruct the history of the Galaxy in its first two billion years and its interaction with other galaxies, shedding new light on enigmatic halo substructures like the Virgo overdensity and Hercules–Aquila cloud. Moreover, a survey of millions of stars will have huge potential for the discovery of kinematically and chemically peculiar stars in as-yet unexplored regions of the Galaxy.

2.7 Complementarity with Other Surveys

While DESI’s spectroscopic survey will by itself yield incisive results in cosmology, its power is increased when combined with other experiments. DESI’s BAO results are directly connected to CMB measurements via its dependence on the acoustic scale, but additional information can be obtained by directly cross-correlating the CMB with the density distribution and redshift space distortions from DESI. Large imaging surveys, including DES and LSST, will provide vast amounts of complementary data, allowing increased precision for both cosmological and neutrino measurements. This combination of imaging and spectroscopic surveys is particularly powerful for distinguishing dark energy from modified gravity models for cosmic acceleration.

2.7.1 Synergies with *Planck* and Future CMB Experiments

The cross-correlation of Planck and potential future CMB experiments with DESI enables cosmological measurements not possible with either individually, and opens up new opportunities to constrain fundamental physics, in the properties of dark energy and gravity discussed in 2.4 and the nature of neutrinos and inflation summarized in 2.5.

On large scales, cross-correlating CMB temperature fluctuations with the galaxy density field measures the Integrated Sachs-Wolfe effect, probing the time evolution of the gravitational potential and independently constraining dark energy [164]. The combination of CMB lensing and the foreground galaxies or quasars will also improve not only the signal-to-noise of CMB lensing leading to stronger cosmological constraints on the matter content, but also our understanding of the foreground tracers in large-scale structure, as lensing allows a clean measurement of the bias of the foreground sources.

The combination of CMB lensing and the RSD measurements from DESI will allow a probe of the two relativistic gravitational potentials independently (see e.g. [55] for an application of this test but for the case of gravitational lensing of background galaxies, not the CMB), testing the GR prediction of their equality over a wide redshift range [165]. CMB lensing and RSD measurements will also provide complementary constraints on the sum and differences of the neutrino masses, that in combination could help infer the neutrino hierarchy.

DESI will provide highly complementary constraints on inflation to those from *Planck* and a number of upcoming CMB small scale temperature and polarization experiments. An exciting realization in inflationary theory is that discerning the scale-dependence, or ‘shape’, of the bispectrum (the 3-point function) could provide a direct insight into the inflationary mechanism, through how non-Gaussianity is generated [166, 167]. CMB 3-point correlation measurements constrain a wide range of primordial bispectrum configurations, while DESI will provide more detailed information about the properties in the squeezed limit, a regime that could provide characteristic information about the underlying mechanism driving inflation e.g. whether it is multi-field, sourced from a non-Bunch Davies vacuum state, or includes non-trivial kinetic terms in the inflationary action.

Cross correlating the galaxy velocity field (inferred from the 3D density distribution) with the CMB will measure the kinetic Sunyaev-Zeldovich (kSZ) effect at the percent level. These measurements provide constraints on more exotic deviations from our standard cosmological models [168, 169, 170]. In addition, these measurements are astrophysically important since the kSZ effect is an unbiased probe of electrons and can be used to inventory the baryons in the Universe [171].

2.7.2 Synergies of DESI with DES and LSST

The massive spectroscopic survey provided by DESI will provide a unique and important complement to direct-imaging science projects currently being planned. We focus here on the Dark

Energy Survey (DES) and the Large Synoptic Survey Telescope (LSST), but DESI will complement other future imaging surveys in similar ways. Although both DES and LSST are located in the Southern Hemisphere, their planned surveys have overlap of a few thousand square degrees with the baseline DESI survey. In addition, only some of the cosmological tests described below rely on overlap between the photometric and spectroscopic surveys.

DESI can provide critical input into photometric redshifts which can help control the systematic uncertainty associated with cosmological measurements from photometric surveys like DES and LSST. For instance, cross correlation of photometric lensing sources with spectroscopic galaxy samples enable the reconstruction of the redshift distribution of the lensing sources [172, 164, 173, 174] providing a critical consistency test on the photometric redshifts used for cosmic shear and/or calibrating the mass of galaxy clusters for cluster abundance tests. Likewise, magnification-based lensing measurements of spectroscopic sources [175] can provide a consistency test for shape systematics and/or photometric redshift systematics in shear-based calibration of cluster masses.

Just as importantly, the combination of photometric and spectroscopic surveys is significantly more powerful than either set of surveys alone. An example is the utility of using galaxy-galaxy lensing, in which one uses the lensing of background galaxies by galaxies from the spectroscopic sample to measure the galaxy-mass cross-correlation of the spectroscopic sample. On small scales, this measures the properties of the host dark matter halo, testing galaxy bias models; on larger scales, it can be used to measure the mass-mass auto-correlation and hence the amplitude of structure [63, 176]. Several studies have forecast cosmological constraints from a combination of DES-like and DESI-like experiments [177, 178, 179, 180], and while the range of assumptions and forecasts varies from work to work, there is agreement that the combination of DES and DESI/LSST gives substantial benefits in terms of measured cosmological and non-cosmological parameters. This is particularly true within the context of modified gravity models, where the combination of surveys enables entirely new types of measurements that are ideally suited for addressing such questions. For instance, recent theoretical work suggests that comparing the shear field generated by galaxy clusters to the corresponding galaxy velocity can significantly improve current modified gravity constraints [181].

As an example of improvement in another type of constraint that can be achieved through the combination of DESI with imaging surveys, Figure 2.16 shows the joint constraint on the sum of the neutrino masses in eV against the dark energy density $\omega_{DE} = \Omega_{DE}h^2$ obtained by combining anticipated results for DESI BAO with LSST weak lensing. Similarly, Figure 2.17 shows prospective constraints in the $\Omega_m - \Omega_\Lambda$ plane obtained by combining anticipated results for DESI BAO with LSST weak lensing (these forecasts assume the surveys are not overlapping on the sky, although it makes practically no difference [84, 182]).

Finally, DES and LSST will provide world-leading samples for supernova cosmology. The BAO and SNe Ia methods for measuring the cosmic distance scale are highly complementary: supernovae excel at low redshifts, where the SNe are brighter and where the BAO is more limited by cosmic variance due to the small cosmic volume. The combination of DESI with ground-based supernovae samples spanning from $z = 0$ to $z \approx 0.8$ will be a powerful view of the distance-redshift relation and the expansion history of the Universe. While we have focused on Figure of Merits drawn only from BAO and the DESI clustering samples, the inclusion of low to intermediate-redshift supernovae provides a notable improvement to current BAO constraints, as highlighted in numerous papers, such as [6, 183]. Essentially one is using BAO to calibrate the relative distance scale provided by the SNe. The redshift overlap of the two methods provides a further systematic cross-check. The exquisite precision of DESI at $z > 0.6$ will find an excellent partner in the DES and LSST supernova samples.

DESI will directly support the coming decade of supernova cosmology by providing spectroscopic




Figure 2.16: Constraint on the sum of the neutrino masses in eV against the dark energy density $\omega_{DE} = \Omega_{DE} h^2$ obtained by combining DESI BAO with LSST weak lensing, in each case including Planck CMB constraints. More powerful constraints are obtained when the full power spectrum from DESI is used. See Table 2.11.

redshifts for many tens of thousands of SNe host galaxies. This will happen both for the faint galaxy survey out to $z \sim 1$, but also with the BGS at $z < 0.4$. Over a 10-year period, a typical (L^*) galaxy has at least a 1% probability of having a detectable SN Ia. This means that the BGS will contain of order 10^5 supernova host galaxies, and the LRG sample of more massive galaxies could produce a comparable number at higher redshift. While photometric redshifts are planned for the large LSST and DES supernova samples, spectroscopic redshifts allow more precision, particularly at low redshift where the uncertainty in the redshift and resulting luminosity distance overwhelm the intrinsic precision of the standard candle. Samples of many tens of thousands of hosts can only be achieved with multi-object wide-field surveys. We note that with DESI there is no need to wait to select the host galaxies after the explosion: at $z < 0.2$, the BGS will include more than half of all SN Ia host galaxies in the survey footprint. Having a pre-existing redshift will also enable better allocation of follow-up resources for rare transients from surveys such as LSST.

2.7.3 Synergies of DESI with *Euclid*/*WFIRST*

Euclid is a medium class European Space Agency survey mission designed to measure Dark Energy [97]. Recently, NASA has become a partner, enabling a group of 40 US astronomers to join the international consortium. *Euclid* will perform a $15,000 \text{ deg}^2$ survey jointly undertaking visible imaging to measure weak lensing and simultaneous near-infrared observations split into sequential imaging (for photometric redshift measurement) and slitless spectroscopy. Two Deep Fields about 2 magnitudes deeper than the wide survey and covering around 20 deg^2 each will be also observed, primarily for calibrations of the wide survey data but also extending the scientific scope of the mission to faint high redshift galaxies, quasars and AGNs. The spectroscopic survey is focused on H α emitting galaxies and is most powerful at high redshifts $1 < z < 2$.

The timeline for DESI is prior to *Euclid* (which is scheduled to launch in 2020), but even in the era of *Euclid*, at redshifts $z < 1$ the combination of LRGs and ELGs that DESI will observe




Figure 2.17: Prospective constraints in the Ω_m – Ω_Λ plane obtained by combining DESI BAO with LSST weak lensing. More powerful constraints are obtained when the full power spectrum from DESI is used. See Table 2.9.

will remain the world-leading data set for spectroscopically confirmed galaxies with good redshift measurements. At $z > 2$ the DESI measurements from Ly- α will also remain unique. Euclid may surpass DESI in the redshift range $1 < z < 2$ provided the slitless spectroscopy is as effective as hoped. DESI could help *Euclid* clustering measurements by providing important information on the potential confusion of the *Euclid* slitless spectroscopy in this redshift range. The combination of Euclid space-based weak lensing with the large spectroscopic samples from DESI will be a strong opportunity for galaxy-galaxy weak lensing, similar to what was discussed in the DES/LSST context in the previous subsection. DESI’s contribution of $z < 1$ lenses is particularly important in this regard.

WFIRST-AFTA is an envisaged NASA mission using a 2.4 m diameter primary mirror satellite being designed to perform a 2000 deg² near-infrared survey, including a slitless spectroscopic component [98]. The current narrow/deep *WFIRST-AFTA* concept is highly complementary to the wide/shallow *Euclid* strategy, and will provide deeper, denser galaxy samples. However, the smaller area covered compared to either *Euclid* or DESI means that the direct expansion rate and growth rate measurements would be weaker.

Comparisons of the precision of the BAO measurement projected for DESI, *Euclid*, and *WFIRST* are shown in Fig. 2.9.

DESI will be highly complementary to the weak lensing surveys to be performed for *Euclid* and *WFIRST-AFTA*, providing spectroscopic galaxy samples at the same redshifts as the matter that is causing the lensing, thus enabling many innovative analyses from these combined datasets. DESI will help in the calibration of photometric redshifts - which are essential for these lensing experiments - and aid in investigating systematic issues such as intrinsic alignments. Likewise, *Euclid* and *WFIRST-AFTA* will greatly enhance the legacy value of DESI, providing high resolution optical and NIR imaging of all DESI targets, greatly improving the prospects for non-dark energy science, e.g., the morphology–density relationship at $z > 1$.

3 Target Selection

The DESI survey will measure with high precision the baryon acoustic feature imprinted on the large-scale structure of the Universe, as well as the distortions of galaxy clustering due to redshift-space effects. To achieve these goals, the survey will make spectroscopic observations of four distinct classes of extragalactic sources – the bright galaxy sample (BGS), luminous red galaxies (LRGs), star-forming emission line galaxies (ELGs), and quasi-stellar objects (QSOs). Each of these categories requires a different set of selection techniques to acquire sufficiently large samples of spectroscopic targets from photometric data. To ensure high efficiency and spectroscopic completeness, we select objects with spectral features expected to produce a reliable redshift determination or a Ly- α forest measurement within the DESI wavelength range.

The characteristics of our baseline samples for each of these target classes are summarized in Table 3.1. This Table specifies the primary redshift range, the photometric bands for targeting, the projected areal density (in terms of number of targets; number of fibers allocated across all pointings accounting for multiple exposures; and the number of useful redshifts resulting per square degree), as well as the total number of objects in the desired class for which redshifts are expected to be obtained for each of these samples. This table may be compared to Table 1 in the Science Requirements Document (SRD). The SRD considers both a threshold survey of 9,000 deg² and a baseline survey of 14,000 deg². Throughout this chapter, we consider only the latter scenario; simulations for reduced focal planes indicate that we would achieve essentially the same sample surface densities as for the baseline scenario, so that sample sizes would simply scale with survey area. In the following sections, we will describe the basis of these numbers in more detail.

Table 3.1: Summary of the properties for each DESI target class. The bands listed are for the target selection, where g , r , and z are optical photometry and $W1$ and $W2$ denote are *WISE* infrared photometry. The exposure densities are increased over the target densities due to some objects being observed on multiple passes. The number of good redshifts and baseline sample sizes (in millions) are for successful redshifts.

Galaxy type	Redshift range	Bands used	Targets per deg ²	Exposures per deg ²	Good z 's per deg ²	Baseline sample
LRG	0.4–1.0	$r, z, W1$	350	580	285	4.0 M
ELG	0.6–1.6	g, r, z	2400	1870	1220	17.1 M
QSO (tracers)	< 2.1	$g, r, z, W1, W2$	170	170	120	1.7 M
QSO (Ly- α)	> 2.1	$g, r, z, W1, W2$	90	250	50	0.7 M
Total in dark time			3010	2870	1675	23.6 M
BGS	0.05–0.4	r	700	700	700	9.8 M
Total in bright time			700	700	700	9.8 M

Summary of Target Samples

The lowest-redshift sample of DESI targets will be the Bright Galaxy Sample (BGS). These galaxies will be observed during the time when the moon is significantly above the horizon, and the sky is too bright to allow efficient observation of fainter targets. Approximately the 10 million brightest galaxies within the DESI footprint will be observed over the course of the survey, sampling the redshift range $0.05 < z < 0.4$ to high density. This sample alone will be ten times larger than the SDSS-I and SDSS-II “main sample” of 1 million bright galaxies observed from 1999-2008.

Above redshift $z = 0.4$, DESI will observe luminous red galaxies (LRGs). These luminous, massive galaxies have long since ceased star formation and therefore exhibit evolved, red composite

spectral energy distributions (SEDs). The BOSS survey targeted these objects to $z \approx 0.6$ using SDSS *gri* colors and measured spectroscopic redshifts using the prominent 4000 Å break continuum feature. While DESI will aim to achieve 350 LRGs/deg² over 14,000 square degrees, the BOSS sample of 119 LRGs/deg² will contribute significantly to our science analyses over the 10,000 deg² footprint in which it exists; DESI may extend this low-redshift sample over a larger footprint, but this is not in the current baseline plan. DESI will target LRGs to $z \approx 1.0$, where they may be most efficiently selected using the prominent 1.6 μm (rest frame) “bump,” which causes a strong correlation between optical/near-infrared (NIR) color and redshift in this regime. We will use 3.4 μm photometry from the space-based Wide-Field Infrared Survey Explorer (*WISE*) to select LRGs efficiently in the redshift range of $0.6 < z < 1.0$. DESI can exploit the 4000 Å break to obtain secure redshifts for LRGs over this full redshift range.

The majority of the spectroscopic redshift measurements for DESI will come from ELGs at redshifts $0.6 < z < 1.6$. These galaxies possess high star formation rates, and therefore exhibit strong emission lines from ionized H II regions around massive stars, as well as SEDs with a relatively blue continuum, which allows their selection from optical *grz*-band photometry. The prominent [O II] λλ3726, 29 doublet in ELG spectra consists of a pair of emission lines separated in wavelength by 2.8 Å. The wavelength ratio of the doublet provides a unique signature, allowing definitive line identification and secure redshift measurements. The goal of the DESI ELG target selection will be to provide a large sample of ELGs with sufficient [O II] line flux to obtain a detection and redshift measurement out to $z = 1.6$.

The highest-redshift target sample will consist of QSOs. We will measure large-scale structure using QSOs as direct tracers of dark matter in the redshift range $0.9 < z < 2.1$. At higher redshifts, we will utilize the foreground neutral-hydrogen absorption systems that make up the Ly-α forest; DESI spectra cover the Ly-α transition at $\lambda = 1216$ Å for objects at $z > 2.1$. We will use optical photometry combined with *WISE* infrared photometry in the W1 and W2 bands to select our primary sample of QSOs. QSOs are ~ 2 mag brighter in the near-infrared at all redshifts compared to stars of similar optical magnitude and color, providing a powerful method for discriminating against contaminating stars. QSOs at $z > 2.1$ used for Ly-α forest measurements do not require homogeneous selection on the sky for cosmological measurements, as we do not rely on the clustering of the QSOs themselves. As a result, DESI may exploit optical variability and additional passbands where available to enhance this sample. Those $z > 2.1$ QSOs which are selected via uniform methods across the sky may also be used to enhance clustering measurements. DESI will obtain additional exposures on the confirmed $z > 2.1$ quasars to measure the Ly-α forest to the required S/N.

Summary of Imaging Required

All DESI target samples will be selected using optical *grz*-band photometry from ground-based telescopes and near-infrared photometry from the *WISE* satellite. The observations assumed in our baseline targeting plan are summarized in Table 3.2. This imaging plan has been developed through a detailed analysis of alternative telescope/instrument combinations. The imaging depths will be at least 24.0, 23.4, 22.5 AB (5σ for an exponential profile $r_3 = 0.45''$) in *g,r,z* and 20.0, 19.3 AB (5σ) in *WISE* W1,W2. All sample magnitude limits quoted in this section are total (model-like) magnitudes for the BGS and for LRGs and ELGs, or PSF magnitudes for QSOs.

The optical imaging for the DESI targets will be provided from three telescopes at two sites, Cerro Tololo and Kitt Peak. The DECam camera on the Blanco 4-m telescope will provide *grz* imaging over 9000 deg² in the DESI footprint at Dec $\leq +30$ deg. The first 6700 deg² of this footprint has been approved as an NOAO survey program during the period August 2014 through July 2017; a proposal to observe the remainder of this footprint is pending. The Bok 2.3-m telescope is providing *gr* imaging over the 5000 deg² region of the North Galactic Cap (NGC) that lies at Dec

$\geq +30$ deg with the existing 90Prime camera. The 220 nights necessary for these observations are guaranteed via an MOU with the University of Arizona / Steward Observatory, and observations began in January 2015. The Mayall 4-m telescope will provide z -band imaging over the same NGC footprint using the existing MOSAIC-2 camera upgraded with 4 red-sensitive CCDs. Those observations will be conducted over 220 nights in January-July 2016 and January-July 2017.

The *WISE* satellite has already obtained and publicly released infrared imaging to sufficient depths for DESI target selection over the full sky. This 13-month survey is being supplemented with an additional 3 years of data taken in the same operational mode from December 2013 which will be used to improve the DESI target selection further. The data from the first 12 months of these continued operations were released on March 26, 2015.

The DESI analyses will be performed separately in each of the three regions of the DESI footprint: the NGC at DEC $> +30$ deg, the NGC at DEC $< +30$ deg, and the South Galactic Cap (SGC). Based on SDSS-III/BOSS experience with separately-calibrated regions, we expect to analyze these separately and combine the cosmological constraints downstream. The DECam and Bok/MOSAIC coverage will have some overlap (at the DEC $\approx +30^\circ$ strip and by targeting specific calibration fields like COSMOS, Boötes, DEEP, etc.) in order to tie together calibrations and understand the subtle variations in target selection resulting from differences in filter+telescope response between the two datasets. The DECam and Bok surveys are now underway and the MOSAIC survey will begin in 2016A.

Table 3.2: Summary of telescopes being used for targeting.

Telescope	Bands	Area deg ²	Location	Status
Blanco DECam	g,r,z	9k	NGC+SGC (Dec $\leq +30$ deg)	Begun 2014B
Bok 90Prime	g,r	5k	NGC (Dec $\geq +30$ deg)	Begun 2015A
Mayall MOSAIC-3	z	5k	NGC (Dec $\geq +30$ deg)	To begin 2016A
WISE-W1	$3.4 \mu\text{m}$	all sky	all-sky	Completed
WISE-W2	$4.6 \mu\text{m}$	all sky	all-sky	Completed

In the remainder of this Section, we demonstrate that our baseline optical/infrared color selections can select the targets listed in Table 3.1, and summarize the key properties of each sample. The accompanying volume of the TDR details the design of the DESI instrument, which informs a spectral simulator presented there. The spectral simulator aids in the design of the targeting strategy (such as magnitude limits), calculates exposure times, and estimates redshift measurement efficiencies. Given the expected target densities and exposure times, the overall survey strategy is developed in Section 4. Included in the survey strategy is an optimized method to tile the sky that maximizes the area covered and number of target redshifts obtained, while minimizing the overall time required for the survey. The outlines for a strategy for fiber allocation are given in the accompanying TDR. This strategy leads to the values given in Table 3.1.

3.1 Targets: Bright Galaxy Sample

3.1.1 Overview of the Sample

The galaxy sample for the BGS will be a flux-limited, r -band selected sample of galaxies. The magnitude limit is determined by the total amount of observing bright time and the exposure times required to achieve our desired redshift efficiency. This target selection is, in essence, a deeper version of the galaxy target selection for the SDSS main galaxy sample (MGS). We explore

the properties of the BGS target sample through mock catalogs created from numerical simulations. These mocks have identical properties to the MGS at low redshift, including the luminosity function, color distribution, and clustering properties. At higher redshifts, the mock BGS is calibrated on data from GAMA ($z \sim 0.3$) and DEEP2 ($z \lesssim 1.0$).

3.1.2 Sample Properties

- *Surface Density:* Figure 3.1 shows the surface density of targets as a function of limiting magnitude. We expect to have a density of just over 800 deg^{-2} for an r -band limit of 19.5, somewhat higher than the goal of $700 \text{ targets per deg}^{-2}$.

- *Redshift Distribution:* Figure 3.2 shows the estimated redshift distribution and space density of galaxies. The upper panel shows the redshift distribution dN/dz in units of 10^3 deg^{-2} per unit redshift. The area under the curve is $800 \text{ targets/deg}^{-2}$. The redshift distribution peaks at $z \sim 0.2$, a factor of 2 higher than the MGS, with a tail out past $z = 0.4$. For comparison, results from GAMA at $r < 19.45$ are shown with the filled circles. The lower panel shows the space density of galaxies in units of comoving $(\text{Mpc}/h)^{-3}$. For reference, the space density of the MGS is shown, as well as the density of BOSS LOWZ+CMASS objects, which is roughly constant at $3 \times 10^{-4} (\text{Mpc}/h)^{-3}$. The BGS sample has a significantly higher density than either the MGS or BOSS out to a redshift of $z = 0.4$. At $z = 0.3$, the sampling of the density field is over an order of magnitude higher in the BGS than in the sum of all SDSS targets.

- *Redshift measurement method:* The broad selection of the BGS will target both star-forming and quiescent galaxies. Redshifts will be obtained from template fits over the full DESI spectral range, with the significance of the fits dominated by the emission lines for star-forming galaxies and by the 4000\AA break and Mg absorption features for quiescent galaxies. Figure 3.3 shows the redshift efficiency as a function of both exposure time and lunar phase for a test sample of galaxies. The test sample is constructed by taking random MGS galaxies and ‘moving’ them further away from the observer by a factor of 2 in redshift. Because the median redshift of the MGS is $z \sim 0.1$, this process creates a test sample with the same median redshift as the BGS sample. We take into account the change in the fraction of light from the galaxy that enters the fiber aperture through redshifting, the change in the angular diameter distance, the change in the point spread function from SDSS to DESI, and the different fiber diameters. `desi_quicksim` is used to create DESI spectra for each test galaxy at a variety of exposure times and lunar phases. Redshifts are obtained using the BOSS redshift code `zfind`, and compared to the true redshift ($2 \times z_{\text{SDSS}}$). Phase, as indicated in the key, is in units of days, with maximum illumination at 14 days and zero illumination at 0 days. Typical BGS observing conditions will be at 10 days, on average. At this phase, the overall redshift success rate is 96% at $t_{\text{exp}} = 6 \text{ min}$, increasing to $\gtrsim 99\%$ at 9 min. Success fraction decreases monotonically with increasing moon illumination. An additional factor in the degree to which the moon affects observations is the angular separation between the moon and the target. All results here are for a separation of 60 degrees.

The results for star-forming and passive galaxies for 10-day phase are shown as well. Galaxies are classified as star forming or passive by their $D_n(4000)$ value, with $D_n(4000) > 1.5$ being passive. At fixed observing conditions, the redshift success rate for star-forming galaxies is lower than for the passive galaxies, indicating that the 4000\AA break is more efficient as a redshift indicator given the spectral noise imparted by the observing conditions. But the redshift success rate for star-forming galaxies is still $\sim 98\%$ for 9 minute exposure times.

- *Large-scale-structure bias:* Estimating the bias of the BGS sample is straightforward due to its completeness in magnitude. We use the abundance matching technique (e.g., [184]) to match galaxies to halos as a function of their luminosity. The bias is then estimated by integrating over




Figure 3.1: The surface density of targets as a function of r -band magnitude. Results are shown for a mock BGS galaxy sample created from numerical simulations. This mock is calibrated to match low-redshift data from SDSS.

the halo mass function, weighted by the number of galaxies per halo. The upper panel in Figure 3.4 shows the bias as a function of redshift obtained with this technique. At low redshift, where the magnitude-limited nature of the survey spans a wide range of absolute magnitudes, the bias is near unity. As redshift increases, the bias monotonically increases. This is for two reasons: for a flux-limited sample, the objects at higher redshift are intrinsically brighter and therefore have higher clustering amplitude, and at higher redshift the bias increases because the amplitude of dark matter clustering is decreasing.

The bottom panel shows the halo masses probed by the BGS target selection as a function of redshift. M_{avg} is the mean halo mass probed. M_{cut} is a cutoff mass scale where halos transition from being too small to host galaxies in the sample to being large enough. Specifically, halos of this mass have a 50% probability of having galaxies in the sample. Significantly above M_{cut} , this probability asymptotes to 100%, but the width of this transition is reflective of the scatter of halo mass at fixed luminosity. This scatter increases with luminosity, which causes M_{avg} to vary more slowly than M_{cut} . For the brightest galaxies, this scatter is so large, M_{avg} is actually below M_{cut} .

- *Target selection efficiency:* The dominant loss of targets is due to fiber assignment inefficiencies. Low-redshift galaxies have higher angular clustering on the sky, which can lead to more contention for fibers in high density regions. However, as described in §4.5, the BGS is being observed in 3 layers to achieve fairly high completeness.

A few percent of galaxies will be lost by deblending errors, superpositions with bright stars, and other artifacts that typically affect imaging catalogs.

- *Areas of risk:* Given the straightforward nature of the target selection, the BGS has minimal risks. There are two possible sources of low-level risk. As shown in Figure 3.3, the redshift efficiency for star-forming objects lags behind that of passive galaxies at fixed observing conditions. The majority of these redshift failures lie in the green valley, in between the main star-forming sequence and the red sequence. These objects have low star formation rates and thus weak emission lines, but do not have stellar populations evolved enough to have strong $D_n(4000)$ values. Dependent on the integration time and observing conditions, the BGS may be incomplete for green-valley objects.

Another possible source of incompleteness is low surface brightness objects, which become more difficult to observe under bright time conditions.




Figure 3.2: *Upper panel:* The redshift distribution of the mock BGS sample. The distribution peaks at $z = 0.18$ with a median redshift of $z = 0.204$. *Lower panel:* The space density of BGS galaxies as a function of redshift. For comparison, the space density of the MGS is shown with the blue curve, and the approximate space density of the full BOSS LRG sample (LOWZ+CMASS) is shown with the dotted line. The space density of the BGS sample is larger than the MGS+BOSS samples up to $z \sim 0.4$.

3.2 Targets: Luminous Red Galaxies

3.2.1 Overview of the Sample

The lowest-redshift coverage for DESI will come from targeting 350 candidate luminous red galaxies (LRGs) per square degree [185]. These objects are both high in luminosity and red at rest-frame optical wavelengths due to their high stellar mass and lack of ongoing star formation. They exhibit strong clustering and a relatively high large-scale-structure bias, which enhances the amplitude of their power spectrum, and hence the BAO signal ([186], [187], [188]). Because of their strong 4000 Å breaks and their well-behaved red spectral energy distributions, low-redshift LRGs at $z < 0.6$ can be selected efficiently and their redshifts estimated based on SDSS-depth photometry [189]. The BOSS survey has targeted 119 LRGs per deg^2 with $z \lesssim 0.6$ using SDSS imaging.

DESI science analyses will incorporate existing BOSS spectroscopic samples (which cover 10,000 deg^2 of the DESI footprint) where available, as well as applying BOSS-like target selection algorithms (in regions not yet covered) to target LRGs at low z . Because the BOSS target selection is well understood and documented in SDSS papers, we will not discuss it further here. Extending the LRG sample to redshifts $z > 0.6$, where the 4000 Å break passes beyond the r band and the optical colors of LRGs overlap with those of red stars, requires different selection techniques, taking advantage of available near-infrared imaging from space. The remainder of this section will focus on the strategy we will use in that domain.




Figure 3.3: The redshift success rate for BGS-like targets. Test targets are created by ‘observing’ MGS galaxies at twice the true redshift of the galaxies. Test spectra are created using `desi_quicksim`, incorporating a lunar model that incorporates the phase, zenith angle of the moon, zenith of the target, and the angle between the target and the moon. Results are shown as a function of exposure time and lunar phase. The green curves show the results for a 10-day lunar phase for passive and star-forming galaxies.

3.2.2 Selection Technique for $z > 0.6$ LRGs

The spectral energy distributions of cool stars exhibit a local maximum around a wavelength of $1.6 \mu\text{m}$, corresponding to a local minimum in the opacity of H^- ions [190]. This feature, commonly referred to as the “ $1.6 \mu\text{m}$ bump”, represents the global peak in the flux density (f_ν) for stellar populations older than about 500 Myr [191], such as those in LRGs. In Figure 3.5 we plot an example LRG template spectrum from [192], illustrating both the strength of this peak and the depth of the 4000 \AA break. The lowest-wavelength channel in *WISE*, the W1 band centered at $3.4 \mu\text{m}$, is nearly optimal for selecting luminous red galaxies; it overlaps the bump at redshift near $z = 1$, so that higher-redshift LRGs will be bright in *WISE* photometry but comparatively faint in the optical. As may be seen in Figure 3.6, a simple cut in $r - W1$ color can therefore select LRGs effectively, while adding in information on $r - z$ color can help in rejecting non-LRGs. *WISE* data are particularly well suited for this application, as the survey depth was designed specifically for detection of L_* red-sequence galaxies to $z = 1$; LRGs are generally significantly brighter than this limit.

We have tested selection techniques using optical *grz* catalogs derived from CFHT Legacy Survey [193] and SDSS Stripe 82 data, NIR imaging catalogs from the *WISE* All-Sky Data Release [194], and redshifts and rest-frame colors derived from DEEP2 spectra [195] or accurate 30-band COSMOS photometric [196] redshifts. A BOSS ancillary program has obtained about 10,000 redshifts of magnitude $z_{SDSS} < 20$ LRG candidates selected using SDSS and *WISE* photometry




Figure 3.4: *Upper panel:* The bias of the BGS sample as a function of redshift. The bias is calculated using the abundance matching model and the space density from Figure 3.2. *Lower panel:* The halo mass scales probed in the BGS sample. M_{avg} is the mean halo mass of the sample. M_{cut} is a cutoff mass scale where halos have 50% probability of containing a galaxy in the sample. The scatter in halo mass at fixed luminosity increases with luminosity, thus uncreases with redshift. This causes the inversion between M_{avg} and M_{cut} when the number density drops below the BOSS value.

with somewhat broader color cuts than DESI will use, which provide additional tests of our basic techniques.

3.2.3 Sample Properties

The baseline LRG selection cuts for DESI are shown by the solid lines in Figure 3.6. This selection, applied to a sample with a total z -band magnitude limit of $z_{AB} = 20.56$, relies on optical photometry in the r and z bands and the infrared photometry in the *WISE* W1 band. DESI target LRGs will often not be detected in the anticipated g band imaging, but are well above the depth limits in the r , z , and W1 bands, having $r < 23$ and $W1 < 19.5$.

This selection is already sufficient to meet all DESI design requirements, though we anticipate further improvements in the future. The major properties of this sample are:

- *Surface Density:* Figure 3.7 shows the effect of changing the limiting magnitude on the surface density of selected targets using the color cuts shown in Figure 3.6. Based upon tests with the public AllWISE catalog and SDSS Stripe 82 Coadd photometry, we find that the baseline sample density of 350 LRG targets/deg² is achieved when selecting objects down to a magnitude limit $z_{AB} = 20.56$. If, instead, we use *Tractor*-based photometry for the full existing *WISE* dataset (cf. Section 3.8), surface densities of objects within the color cut are about 20% higher (due primarily to the mitigation of confusion effects), allowing our target surface density to be achieved with a




Figure 3.5: A template spectrum based upon observations of the nearby elliptical galaxy NGC 4552, drawn from the work of [192]. The spectrum f_ν is plotted as a function of rest-frame wavelength; we overplot the total (telescope + instrument + detector) response curves for DECam *grz* and *WISE* W1 and W2 imaging at the appropriate rest frame wavelengths for an LRG at $z = 0.9$. The 1.6 micron bump, the key spectral feature that enables our LRG selection method, corresponds to the peak in this spectrum. In the inset, we plot flux f_λ over a limited wavelength range in order to illustrate clearly the 4000 Å break and the abundance of spectral absorption features in this vicinity, which will be exploited by DESI to measure redshifts for LRGs.

magnitude limit of $z_{AB} = 20.45$ instead. Tests with COSMOS photometric redshifts indicate that the objects in our selection box gained from the *Tractor* analysis are all both red and luminous; only a few $z_{AB} < 20.45$ LRGs in this field are missed by our selection, but they are generally both at the low-redshift and blue (and hence less strongly biased) end of the sample.

Based on the results of the BOSS ancillary *WISE* LRG program, we expect high ($> 98\%$) redshift completeness for $z_{AB} < 20$ LRGs with one DESI visit, for $z_{AB} < 20.38$ with two visits, or for $z_{AB} < 20.57$ with three visits. For our baseline sample, a mean of two visits per object will be required (given the fractions of the sample with $z_{AB} < 20$ or $z_{AB} > 20.38$). The baseline sample is roughly as large as feasible while simultaneously rejecting $z < 0.6$ objects and requiring an average of at most two observations per LRG. Given their high large-scale-structure bias, fainter LRGs may be more valuable than ELGs at the same redshift, and hence may be worth the longer exposure times necessary; more work on optimization remains to be done. Out of the DESI LRG target sample, about 50 per square degree will have been observed by eBOSS where the footprints overlap, proportionately reducing the final number of LRG targets in these regions.

- *Redshift Distribution:* We have estimated the redshift distributions resulting from the DESI baseline target selection (see Figure 3.8) by using COSMOS photometric redshifts or spectroscopic redshifts from our SDSS-III/BOSS ancillary program. Specifically, for the latter we applied the DESI selection cuts to SDSS Stripe 82 + *WISE* photometry, and then assigned the selected galaxies the spectroscopic redshift of the nearest-color object from our BOSS ancillary program. The larger noise in the SDSS imaging over the ancillary program’s footprint causes the selection to pick up lower-redshift objects, while the high-redshift tail is suppressed by the lack of redshifts at $20 < z_{\text{SDSS}} < 20.56$, making the resulting redshift distribution somewhat lower than DESI should




Figure 3.6: An optical/near-infrared color-color diagram for galaxies observed by both the CFHT Legacy Survey and WISE in the COSMOS field, where highly accurate 30-band photometric redshifts are available and used to label points here. In this and subsequent figures, r indicates SDSS r -band magnitude, z indicates SDSS z_{AB} , and W1 indicates *WISE* 3.4 μm AB magnitude. Galaxies with LRG-like spectral energy distribution also having $z > 0.6$ are indicated by points color-coded according to their redshift, whereas small black points indicate blue galaxies at all redshifts. The dashed lines indicate the borders of our LRG selection box; our baseline sample assumes that objects above and to the right of these lines that also have magnitude $z_{AB} < 20.56$ will be targeted by DESI as high-redshift LRGs.

attain. In contrast, the COSMOS imaging is more comparable in depth to that we anticipate, and photo- z 's are available to much fainter than $z = 20.56$, but due to the small area of the field sample/cosmic variance yields strong fluctuations in the redshift distribution. Even given these limitations, we find that our sample selection meets or exceeds all requirements for the DESI baseline LRG sample.

As this figure shows, we have a particularly large density of objects at $z < 0.8$ and will likely down-sample at those redshifts accordingly (e.g., by using a brighter magnitude limit for objects with blue $r - z$ colors). The apparent magnitude of LRGs is strongly correlated with their redshift, allowing us to sculpt the LRG redshift distribution efficiently.

- *Redshift measurement method:* LRGs exhibit a prominent break in their spectral energy distribution around 4000 Å (rest-frame), associated with multiple strong absorption-line features. This feature will be covered by the DESI spectrograph at redshifts up to $z = 1.45$. Our exposure times per target are set to achieve equivalent signal-to-noise at the wavelengths of interest as our BOSS ancillary program targeting $z_{SDSS} < 20$, $z > 0.6$ LRGs attained in one hour of SDSS exposure time. We therefore expect to obtain highly-secure redshifts for a comparable fraction of targets ($> 98\%$) as in that ancillary program.

- *Large-scale-structure bias:* In order to predict the strength of the BAO feature in galaxy clustering measurements, we must assume a value for the ratio of galaxy clustering to dark matter clustering, commonly referred to as the large-scale structure bias. On large scales this may be approximated as a function of redshift that is independent of scale, $b(z)$. We can anticipate that the bias for $z > 0.6$ luminous red galaxies should be at least as large as that of BOSS LRGs, as only




Figure 3.7: Surface densities of targeted candidate $z > 0.6$ LRGs as a function of limiting z -band magnitude. We plot here the surface density of objects that lie within the target selection box shown in Figure 3.6 as a function of their z_{SDSS} magnitude, as determined from data in SDSS Stripe 82. We also indicate our goal density of 350 targets per square degree via the magenta dashed line. Our baseline LRG sample size is attained at a depth of $z_{AB} < 20.56$ for the AllWISE catalog or $z_{AB} < 20.45$ for *Tractor*-based catalogs. At the fainter of these limits, an average of two spectroscopic measurements per LRG will be required to attain secure redshifts for $> 98\%$ of targets.

the most extreme objects will be able to assemble a large amount of mass and cease star formation by this earlier epoch. We therefore assume a bias of the form $b(z) = 1.7/D(z)$, where $D(z)$ is the growth factor; this matches the value measured by SDSS-I at $z = 0.34$ [186] and by SDSS-III at $z = 0.57$ [197]. We have extrapolated this trend to $z = 1$ for the DESI LRGs.

- *Target selection efficiency:* Targets selected as LRGs could fall short in several ways: they could fail to yield redshifts entirely; they could prove to be stars rather than galaxies; they could be outside of the desired redshift range; or they could turn out to be blue (i.e., star forming and less highly biased). Based on results from the BOSS ancillary program, we expect to obtain redshifts for $> 98\%$ of LRGs targets, as described above. Roughly 2% of the objects targeted via the baseline selection box (which could be further optimized) are stars. 98% of the galaxies selected are at $z > 0.6$, while 98% of the galaxies selected prove to have red-sequence rest frame colors. If we treat all failure modes as independent (the worst-case scenario), this yields a net target selection efficiency of 92%; i.e., more than 92% of all DESI LRG targets will be luminous red galaxies in the correct redshift range with a secure redshift measurements.

- *Areas of risk:* There are few sources of risk in our LRG selection, the most important of which is the possibility that the COSMOS field is unrepresentative of the overall survey and instead contains (due to Poisson statistics or cosmic variance) an unusually large fraction of galaxies with red colors at $z > 0.6$. Although the likelihood of this is very low, at worst, this could degrade the target selection efficiency to near 90%. The second potential source of risk is that the redshift success rate for LRGs is not simply a function of the signal-to-noise ratio, in which case we can not map our BOSS ancillary experience to DESI. The best way to retire both of these risks will be via spectroscopic observations of a straw-man DESI LRG sample with a red-sensitive, wide-field, multi-object spectrograph well in advance of the DESI survey.




Figure 3.8: DESI LRG redshift distribution for our candidate sample from two studies: (*black*) Photometric redshift distribution from the CFHT-LS imaging in the COSMOS field, which has accurate photometry and full redshift coverage but suffers from high sample variance (as seen from the feature at $z \approx 0.77$). (*blue*) Spectroscopic redshift distribution for galaxies selected from the SDSS Stripe 82 photometry and assigned the redshift to objects with the same color from a BOSS ancillary program. The latter sample has low sample variance, but the high-redshift tail is suppressed by the lack of redshifts at $20 < z_{\text{SDSS}} < 20.56$. Shown in red is the redshift distribution of low- z LRGs, many already observed by SDSS-I/II and SDSS-III/BOSS, which will be included in the DESI analysis.

To summarize, the luminous red galaxy selection methods used for our baseline plan will yield a high-bias sample of about 315 LRGs/deg² (assuming 90% efficiency net) out of a sample of 350 targets/deg²; almost all will be galaxies at $z > 0.6$. To be conservative, our projections assume that only 86% of the targeted LRGs (i.e., 300 per square degree) will in fact be $z > 0.6$ luminous red galaxies. Combined with BOSS LRGs at lower redshift, this will allow us to measure the BAO scale from $0 < z < 1$. This sample allows direct comparisons to cosmological results provided by the ELG sample in overlapping redshift ranges, providing a key test for systematic effects.

3.3 Targets: Emission Line Galaxies

3.3.1 Overview of the sample

Emission-line galaxies (ELGs) constitute the largest sample of objects that DESI will observe. Such objects exhibit strong nebular emission lines originating in the ionized (“H II”) regions surrounding short-lived but luminous massive stars. ELGs are typically late-type spiral and irregular galaxies, although any galaxy that is actively forming new stars at a sufficiently high rate will qualify as an ELG. Because of their vigorous ongoing star formation, the integrated rest-frame colors of ELGs are dominated by massive stars, and hence will typically be bluer than galaxies with evolved stellar populations such as LRGs. The optical colors of ELGs at a given redshift will span a larger range than LRGs due to the much greater diversity of their star formation histories and dust properties.

DESI leverages the fact that the cosmic star formation rate was roughly an order of




Figure 3.9: Example rest-frame spectrum of an ELG showing the blue stellar continuum, the prominent Balmer break, and the numerous strong nebular emission lines. The inset shows a zoomed-in view of the [O II] doublet, which DESI is designed to resolve over the full redshift range of interest, $0.6 < z < 1.6$. The figure also shows the portion of the rest-frame spectrum the DECam *grz* optical filters would sample for such an object at redshift $z = 1$.

magnitude higher at $z \sim 1$ than today, which causes galaxies with strong line emission to be very common at that epoch [198, 199, 200]. Fig. 3.9 shows an example rest-frame spectrum of an ELG, which is characterized by a blue stellar continuum dominated by massive stars, a Balmer break at $\sim 3700 \text{ \AA}$ (whose strength depends on the age of the stellar population), and numerous nebular emission lines, the most prominent of which are $H\alpha$ $\lambda 6563$, $H\beta$ $\lambda 4861$, the higher-order Balmer lines, and the forbidden [O III] $\lambda\lambda 4959, 5007$ and [O II] $\lambda\lambda 3726, 3729$ nebular emission-line doublets. The inset provides a zoomed-in view of the [O II] doublet (assuming an intrinsic line-width of 70 km s^{-1}), which the DESI instrument is designed to resolve over the full redshift range, $0.6 < z < 1.6$. By resolving the [O II] doublet, DESI will avoid the ambiguity of lower-resolution spectroscopic observations, which cannot differentiate between this doublet and other single emission lines [201].

3.3.2 Selection Technique for $z > 0.6$ ELGs

The DESI/ELG targeting strategy builds upon the tremendous success of the DEEP2 galaxy redshift survey, which used cuts in optical color-color space to effectively isolate the population of $z \gtrsim 0.7$ galaxies for follow-up high-resolution spectroscopy using the Keck/DEIMOS spectrograph [203, 195]. More recently, as part of an approved SDSS-III ancillary program, [204] have confirmed that optical color-selection techniques can be used to optimally select bright ELGs at $0.6 < z < 1.7$.

In Fig. 3.10 we plot the $g - r$ vs $r - z$ color-color diagram for those galaxies with both highly-secure spectroscopic redshifts and well-measured [O II] emission-line strengths from the DEEP2 survey of the Extended Groth Strip (EGS) [195]. The *grz* photometry of these objects is drawn from CFHTLS-Deep observations of this field [202], degraded to the anticipated depth of our DECam imaging (see §3.6.1). As discussed in the next section, we expect to achieve a very high redshift success rate for ELGs with integrated [O II] emission-line




Figure 3.10: Optical $g-r$ vs. $r-z$ color-color diagram based on spectroscopy from the DEEP2 Galaxy Redshift Survey and imaging from CFHT LS [202], illustrating our strawman selection for ELGs at $z > 0.6$ with significant [O II] emission-line flux. This plot shows that strongly [O II]-emitting galaxies at $z > 0.6$ (blue points) are in general well-separated from both the population of lower-redshift galaxies (pink diamonds) and from the locus of stars in this color space (grey contours). The strawman selection box (thick black polygon) has been designed to optimize the number of strongly [O II]-emitting ELGs at $z \sim 1$ selected while simultaneously minimizing contamination from stars and lower-redshift interlopers.

strengths in excess of approximately $8 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$.⁴

Fig. 3.10 shows that strongly [O II]-emitting galaxies at $z > 0.6$ (blue points) are well isolated from the population of lower-redshift galaxies (pink diamonds), as well as from the stellar locus (grey contours), except at the very bluest colors (as the most highly star-forming galaxies have relatively flat spectra with little Balmer break, they exhibit similar colors at all redshifts). The separation between galaxies above and below $z \simeq 0.6$ occurs due to the spectrum blueward of the Balmer break ($\lambda_{\text{rest}} \sim 3700 \text{ \AA}$; cf. Figure 3.9) shifting into the r -band filter, which rapidly reddens the $r-z$ color; similarly, at $z \gtrsim 1.2$ the Balmer break moves into the z -band filter, causing both the $g-r$ and $r-z$ colors to be relatively blue at higher redshifts. The black polygon in Figure 3.10 delineates the candidate selection box we have used to isolate the population of strongly [O II]-emitting, intermediate-redshift ELGs. By targeting galaxies in this box to a depth of $r_{AB} = 23.4$, we strike a balance between maximizing the number of $z \sim 1$ ELGs with significant [O II] flux targeted, while simultaneously minimizing contamination from stars and lower-redshift galaxies.

3.3.3 Sample Properties

The baseline ELG selection criteria for DESI are based on our analysis of the DEEP2/EGS survey data, which targeted galaxies more than half a magnitude fainter and with consider-

⁴This integrated [O II] flux corresponds to a limiting star-formation rate of approximately 1.5, 5, and $15 M_{\odot} \text{ yr}^{-1}$ at $z \sim 0.6$, 1, and 1.6, respectively, which lies below the ‘knee’ of the star formation rate function of galaxies at these redshifts [205, 206].




Figure 3.11: Surface density of ELGs within our candidate color selection box as a function of limiting r -band magnitude. The solid black line indicates the surface density of objects that lie within the target selection box shown in Figure 3.10 as a function of their r_{AB} magnitude, as determined from CFHTLS-Deep data [193]. The blue dashed line shows the surface density of selected targets when all measurements are degraded to match the expected errors from the DECam Legacy imaging. The dashed magenta line shows our goal density of 2400 targets deg^{-2} , which is achieved at a depth of $r_{AB} \lesssim 23.4$ in either scenario.

ably higher spectroscopic signal-to-noise ratio than DESI. Because of this greater depth, we anticipate that any galaxies with sufficiently strong [O II] flux to yield a redshift with DESI will also yield a successful redshift measurement in DEEP2. We have also cross-verified our selection criteria and redshift distributions for ELGs using data from the 1.3 deg^2 COSMOS field [207] and from the 0.6 deg^2 VVDS-Deep field [208]; both of these samples give consistent results, well within the expected variation due to both sample variance and systematic differences between the samples. Our selection, when applied to imaging with magnitude limits of $g_{AB} = 24$, $r_{AB} = 23.4$ and $z_{AB} = 22.5$ (i.e., the anticipated depth of DECam Legacy imaging), is sufficient to meet all DESI science requirements, though we anticipate further improvements in the future. The major properties of this sample are:

- *Surface Density:* In Fig. 3.11 we show the surface density of candidate ELGs in our grz selection box (see Fig. 3.10) as a function of the r -band magnitude limit. At a depth of $r_{AB} = 23.4$, we achieve our goal of 2400 targets per square degree. As we discuss below, we conservatively estimate that at least 65% of these will be bona fide ELGs in the redshift range $0.6 < z < 1.6$ with a strong enough [O II] emission-line doublet (in tandem with other nebular emission lines available at $z \lesssim 1$) to yield a secure redshift. Out of this sample, at most 270,000 ELGs over $500 - 1,500$ square degrees may be targeted by the SDSS-IV/eBOSS survey, representing a very small fraction of the overall DESI targets.

- *Redshift Distribution:* Fig. 3.12 shows the anticipated redshift distribution of our candidate grz -selected sample of ELGs, determined based on those DEEP2/EGS objects which are both selected by our candidate cuts (after degradation to DECam Legacy errors) and exhibit sufficient [O II] flux for DESI redshift measurements to succeed, reweighted to account




Figure 3.12: Expected redshift distribution of ELG targets based on our analysis of the DEEP2/EGS survey data (see Figure 3.10). The overall normalization of the distribution has been fixed to $1280 \text{ ELGs deg}^{-2}$ (from a targeted sample of $2400 \text{ targets deg}^{-2}$) to reflect conservative estimates of the overall efficiencies of fiber assignment, target selection, and redshift measurement. The ELG redshift distribution drops to a level where shot noise dominates errors in BAO measurements (i.e., $\bar{n}P < 1$) only at $z \gtrsim 1.3$ (dashed blue line).

for DEEP2 target selection rates.⁵

The dashed blue line in the figure shows the surface density at a given redshift for which $\bar{n}P = 1$ (when evaluated at wave number $k = 0.14 \text{ h Mpc}^{-1}$ and orientation relative to the line-of-sight $\mu = 0.6$); below this limit, shot noise will dominate errors in measuring the BAO signal (cf. §2.4.2). Our candidate ELG selection exceeds the $\bar{n}P = 1$ curve to redshift $z \sim 1.3$.

- *Redshift measurement method:* The adopted *grz* color-cuts are designed to maximize the selection of galaxies at $z \sim 1$ with significant [O II] emission-line flux. In Fig. 3.13 we plot [O II] flux as a function of redshift using the DEEP2/EGS sample. The pink curve shows the limiting [O II] flux above which DESI simulations predict that we will detect multiple emission features at $> 7\sigma$ (cf. Performance section of accompanying TDR); in such cases, we expect to measure a secure redshift at least 95% of the time. For galaxies at $z \lesssim 1$ the redshift will be constrained using [O II] as well as the [O III] doublet and $H\beta$, while beyond $z \gtrsim 1$, the [O II] doublet will be the only strong lines present in the observed-frame optical spectra.

- *Large-Scale Structure Bias:* We may estimate the linear clustering bias of our sample of ELGs relative to their dark matter halos using the DEEP2 data. Employing methods similar to those of [209] and [210], we have measured the clustering of ELGs (selected using

⁵DEEP2 does not cover [O II] at $z < \sim 0.8$ or $z > \sim 1.4$. We handle this at low redshifts by assigning [O II] fluxes from galaxies of matching color and luminosity at slightly higher z to those objects lacking [O II] coverage in DEEP2. For $z > 1.4$, we plot a power-law extrapolation of the redshift distribution measured at lower redshift, as DEEP2 would in general not obtain a redshift at all for objects where [O II] is past the red end of the spectrum. An analysis of COSMOS photometric redshifts for objects meeting our selection cuts suggests that this extrapolation if anything underestimates the number of objects at $1.4 < z < 1.6$.




Figure 3.13: [O II] flux as a function of redshift for DEEP2/EGS galaxies. The light blue squares represent all galaxies in the sample, while the dark blue points are those objects meeting our ELG selection criteria (see Fig. 3.10). The pink curve shows the limiting [O II] flux above which simulations indicate that DESI will detect multiple emission features (where the [O II] doublet constitutes two features) at $> 7\sigma$. With multiple lines detectable at such high significance, we conservatively expect to obtain a secure redshift at least 95% of the time (cf. Performance section of accompanying TDR).

an earlier version of our current cuts) at quasilinear scales of $1 - 10 h^{-1}$ Mpc in three overlapping redshift bins centered at $z = 0.87, 1.0$ and 1.2 . The observed clustering is consistent within errors at all redshifts, even though the amplitude of matter clustering was lower at higher z [211]. The observations can thus be described by a galaxy bias which is inversely proportional to the growth factor of dark matter fluctuations; based upon our measurements we adopt $b(z) = 0.84/D(z)$, where $D(z)$ is the growth factor at redshift z ($D(z) = 1$ today). This increase in the bias for star-forming galaxies as redshift increases is consistent with other studies of similar objects at $z=0.5-2.2$ [212, 213, 214].

- *Target selection efficiency:* Targets selected as ELGs could fall short in several ways: they could entirely fail to yield a redshift (e.g., if the galaxy is at $z \gtrsim 1.63$ then no strong emission lines will be detected by DESI); they could prove to be low-redshift galaxies, $z < 0.6$; they could be QSOs instead of galaxies (and hence useful for higher-redshift clustering analyses but likely outside the redshift range of the ELGs); or they could be stars. Based on the DEEP2/EGS sample, we estimate that $\sim 10\%$ of the objects targeted via the baseline selection criteria are expected to be stars, $\sim 5\%$ will be lower-redshift interlopers, and $\sim 5\%$ will be at $z \gtrsim 1.6$, while contamination from QSOs is expected to be negligible. Combining all these factors, the fraction of ELG targets that are in fact galaxies in the correct redshift range is approximately 80%. Among these objects, about 85% will have a large enough [O II] flux to securely measure a redshift more than 95% of the time (see Fig. 3.13). Combining all these factors with the 78% fiber assignment rate expected for an input target density of $2400 \text{ targets deg}^{-2}$ (see the section on fiber assignment in the accompanying TDR), we obtain an a final density of $1220 \text{ ELGs deg}^{-2}$.

- *Areas of risk:* The primary source of risk in our ELG selection is the limitations of

the datasets available for developing and assessing selection algorithms. DEEP2 is the only large current survey which resolves the [O II] doublet critical for obtaining secure redshifts at $z > 1$; however, due to the $z > 0.75$ color cut applied by DEEP2 in three of four survey fields, it can be used to assess the low-redshift tail of the ELG selection in only a limited area, the Extended Groth Strip used for all analyses here. Because of the limited area, the number of DEEP2 ELGs within our color box is small, so both Poisson noise and sample/cosmic variance have a significant effect on predicted redshift distributions. Furthermore, the lack of DEEP2 coverage of [O II] at $z > \sim 1.4$ means that our assessments of performance in that regime are subject to some amount of uncertainty; however, with any other extant dataset, even more assumptions and extrapolations would be necessary. Because of this, projections of DESI ELG performance based upon the available data could prove inaccurate at some level. The consistency of VVDS and COSMOS results with the DEEP2-based predictions builds confidence that these inaccuracies are not substantial.

The second potential source of risk which would cause performance to fall short of our projections is that the redshift success rate for DESI ELGs could not simply be a function of signal-to-noise ratio, but may also depend in more subtle ways upon the instrumental resolution and the intrinsic galaxy velocity dispersions. For example, it would be difficult to directly discriminate between [O II] or another single-line feature at lower redshift for a population of ELGs with unusually large velocity dispersions $\sigma_v > 150 \text{ km s}^{-1}$ (though the rarity of low-luminosity objects with extremely high velocity dispersions, as would be implied by a false identification, may allow such cases to be resolved). The best way to retire all of these risks is to perform spectroscopic observations of straw-man DESI ELG samples using a spectrograph having similar spectral resolution to DESI, and covering the full optical wavelength window.

To conclude, the ELG selection methods used for our baseline plan will yield 2400 targets deg^{-2} . From these targets, DESI should securely measure redshifts for approximately 1220 ELGs deg^{-2} in the redshift range $0.6 < z < 1.6$ (see Table 3.1). This sample will enable constraints on cosmological parameters over a broad redshift range centered on $z \sim 1$, which can be directly compared to results from the independently observed samples of LRGs at $z < 1$ and quasars at $z > 1$.

3.4 Targets: QSOs

3.4.1 Overview of the sample

The highest-redshift coverage of DESI will come from quasars (a.k.a. quasi-stellar objects, or QSOs), extremely luminous extragalactic sources associated with active galactic nuclei. QSOs are fueled by gravitational accretion onto supermassive black holes at the centers of these galaxies. The QSO emission can outshine that of the host galaxy by a large factor. Even in the nearest QSOs, the emitting regions are too small to be resolved, so QSOs will generally appear as point sources in images. These are the brightest population of astrophysical targets with a useful target density at redshifts $z > 1$ where the population peaks [215, 216].

DESI will use QSOs as point tracers of the matter clustering mostly at redshifts lower than 2.1, in addition to using QSOs at higher redshift as backlights for clustering in the Ly- α forest. This enlarges the role of QSOs relative to the BOSS project, which only selected




Figure 3.14: Cumulative surface density of quasars (objects per deg^2) as a function of g magnitude for $z > 0.9$, derived from different estimates of the QSO luminosity function.

QSOs at $z > 2.15$ for use in the Ly- α forest, and enhances their role relative to eBOSS where QSOs are used in a similar fashion as in DESI although with lower densities. DESI will select 170 QSOs per deg^2 over its footprint, of which 50 per deg^2 will be at $z > 2.1$ and suitable for the Ly- α forest.

DESI pilot programs have answered the long-standing uncertainties in the faint end of the QSO luminosity function [217, 216]. The surface density for $z > 0.9$ QSOs derived from these studies is shown in Figure 3.14, along with previous estimates from [218] (15% lower) or from the LSST science book [219, 220] (60% higher). Brighter than magnitude $g = 23.0$, we infer that a *complete* QSO sample would contain about 160 QSOs per deg^2 at $z < 2.1$ and about 68 at $z > 2.1$. DESI will target and obtain redshifts for 120 and 50 QSOs per deg^2 in the redshift ranges $z < 2.1$ and $z > 2.1$, respectively.

Because of their point-like morphologies, and with *photometric* characteristics that mimic faint blue stars in optical wavelengths (Figure 3.16, middle plot), QSO selection is challenging. The photometric selection used by BOSS to target Ly- α QSOs at $z > 2.15$ has attained a 42% targeting efficiency (i.e., fraction of targets that prove to have the desired class and be in the desired redshift range), yielding 17 $z > 2.15$ QSOs per deg^2 down to the SDSS photometric limit of $g < 22.1$ [11]. The selection technique for DESI needs to achieve a minimum efficiency of about 70%; unlike for BOSS, however, QSOs at $z < 2.15$ would be considered successes. A baseline scheme for QSO selection that already achieves our goals for DESI is presented below.

3.4.2 Selection Technique

QSOs commonly exhibit hard spectra in the X-ray wavelength regime, bright Ly- α emission in the rest-frame UV, and a power-law spectrum behaving as $F_\nu \propto \nu^\alpha$ with $\alpha < 0$ in the mid-infrared bands [221] (c.f. Figure 3.15). In the mid-optical colors, QSOs at most redshifts are not easily distinguished from the much more numerous stars. Successful selection of a highly-complete and pure QSO sample must make use of either UV or infrared photometry; DESI relies upon optical and infrared photometry for its baseline selection.

The QSO target selection is a combination of optical-only and optical+IR selections.




Figure 3.15: QSO spectrum exhibiting the main emission lines used in their identification.

The greatest separation from the stellar locus in the optical comes from ugr colors where the “UV excess” in $u - g$ produces bluer colors than those of most stars (Figure 3.16 left). In the absence of u band in the baseline imaging plan, the bulk of the QSO targets are identified in an optical+IR selection, where the excess infrared emission from QSOs results in a clear segregation from stars with similar optical fluxes. Stellar SEDs indeed sample the rapidly declining tail of the blackbody spectrum at those wavelengths, where QSOs have a much flatter SED. A color selection is defined by cuts in $g - r$ and $r - W1$ shown in Figure 3.16c. The *WISE* data are available on the whole sky, and are photometered deeper than the public *WISE* catalogs using the Tractor-forced photometry (see section 3.8). Although *WISE* and optical data are not synchronous, the color difference between QSOs and stars is so large that QSO variability has minimal effect on the color selection. The *WISE* satellite has been reactivated, and will improve by a factor of two in signal-to-noise prior to DESI. A supplemental optical-only selection increases the completeness of the DESI QSOs at the faintest magnitudes, although this may not be necessary once the deeper *WISE* data is in hand.

This baseline QSO target selection was tested in the deep CFHT D3 field where we led a DESI pilot survey (as an ancillary program in BOSS) in order to build a catalog of spectroscopically identified QSOs at all redshift. This pilot survey aimed at an optimal selection of $g < 22.5$ QSOs from combined color and variability information, using a large $ugriz$ and near-infrared data set. Our baseline selection was then done using optical grz catalogs derived from CFHT Legacy Survey [193] combined with near-infrared photometry from *WISE*. The primary selection in $g - r$, $r - W1$ space is shown in the right-hand panel of Figure 3.16, selecting to depths $r = 23.0$. This is supplemented by a tight optical color-box selection, restricted to the region bounded by the wedge at high $g - z$ and low $g - r$ in the central panel of Figure 3.16. This second selection adds few candidates compared to our primary selection, but increases our efficiency in particular for faint events that cannot have a reliable IR flux measurement with current *WISE* data. In both cases, we restrain the selection to objects with point-like morphology. The union of these selections results in about 170 QSOs per deg^2 , with 50 per deg^2 at $z > 2.1$, and a contamination of about 80 stars per deg^2 .

DESI may supplement its high-redshift QSOs with more sophisticated selection algorithms and other supplementary photometry as it becomes available. Time-domain data en-

able variability selection methods (as described in Section 3.4.5). UV (u -band) data improve QSO selection, and allow discrimination between low-redshift and high-redshift QSOs. Algorithmically, neural-network based algorithms [222] and an extreme deconvolution method that models the distributions of stars and quasars at the flux limit [223] have been in use by BOSS where they allowed an increase of up to 20% in selection efficiency over traditional selection algorithms [224]. They are also applied, and thus further tested, in eBOSS. A combination of these additional data and algorithms will allow DESI to target high-redshift QSOs in excess to those currently planned, with a small impact on the overall fiber budget.

The main contaminants to a grz +*WISE* QSO selection are very low-redshift star-forming galaxies with strong PAH emission, currently excluded using a star-galaxy separation based on ground-based optical imaging; a few high-redshift obscured galaxies, which are rare at bright optical magnitudes; and faint stars that artificially drift into the QSO locus because of poor optical photometry.

3.4.3 Sample Properties

Selection using optical grz and near-infrared data for the deep CFHT D3 field achieved a performance at the level of our goals for the DESI sample. Application of additional data and more sophisticated selection algorithms may be used to boost the high-redshift QSO densities. The major properties of the baseline DESI QSO sample are :

- *Surface Density:* The grz +*WISE* color-box selection yields a total of 255 targets per deg^2 to a limit $g = 23$, of which about 110 per deg^2 are confirmed QSOs with $z < 2.1$ and about 50 per deg^2 are QSOs at $z > 2.1$, similar to the required densities of Table 3.1. Based on the QSO luminosity function of [216], this corresponds to about 70% of all QSOs in this magnitude range. We anticipate that the deeper *WISE* data expected before the start of DESI will give us the opportunity to improve the selection by increasing the completeness and decreasing the stellar contamination even further.




Figure 3.16: The ugr and near-infrared (*WISE* W1 band) colors of objects photometrically classified as stellar point-like objects (blue points) and those spectroscopically classified as QSOs, with green contours indicating tracer QSOs at $z < 2.1$ and red contours for Ly- α QSOs with $z > 2.1$. Note the large discrimination between QSOs and stars in the right-hand panel. Black lines mark the boundaries of the selection regions described in the text.




Figure 3.17: Expected distribution of QSO redshifts from DESI. The expected redshift distribution of the QSO sample is indicated by the thick red histogram using the DESI targeting efficiency of 70% at $g < 23.0$. This is in excellent agreement with the DESI pilot survey in the CFHT LS W3 region (blue points). For comparison are the QSO luminosity function to $g < 22.5$ (blue dashed line) and $g < 23.0$ (red dashed line).

- *Redshift distribution:* The expected redshift distribution of the QSO sample is illustrated in Figure 3.17 as the thick red histogram, which is based upon assuming that 70% of all QSOs brighter than $g < 23$ will be included in our sample. For comparison, we show on the same plot the QSO density achieved with the DESI pilot survey (see description in section 3.4.2) in the CFHT LS W3 region (blue points), as well as the QSO luminosity function to $g < 22.5$ (blue dashed line) and $g < 23$ (red dashed line). At $z > 1$, the completeness of our pilot sample approaches 100% uniformly over all redshifts. The small loss that appears at $z < 0.9$ is due to the selection of stellar-like objects only where extended light from a QSO’s host galaxy can make them appear morphologically extended.

- *Redshift measurement method:* The key features contributing to the classification and redshifts of QSOs are the Ly- α , CIV, CIII] and MgII emissions (c.f. Figure 3.15). From our experience during BOSS and with pilot programs (with BOSS and larger telescopes such as the MMT), we estimate that in a single DESI visit we will fail to obtain redshifts for QSO targets about 14% of the time, mostly for objects at $g > 22.5$ [216]. All QSO targets will be observed early in the survey. Those confirmed to be QSOs at $z > 2.1$ will be re-observed in subsequent passes over the sky in order to obtain higher signal-to-noise spectra of the Ly- α .

- *Large-scale-structure bias:* QSO bias has been measured in BOSS via QSO-Ly- α cross-correlation studies to be about 3.6 at $z = 2.4$ [225], in agreement with previous measurements [226, 227]. For QSOs at lower redshifts, we project a bias of the form $b(z) = 1.2/D(z)$, where $D(z)$ is the growth factor. At $z > 2.1$, clustering information is computed from the transmitted flux in the Ly- α forest and not directly from correlations between objects; the flux bias of Ly- α absorbers is estimated to be about -0.2 (it is negative because a larger matter density implies a higher absorption and thus a lesser transmitted flux) [228], and is strongly enhanced along the line of sight by redshift-space distortions.

- *Target selection efficiency:* From the first pass of targeting over the sky, we expect to identify 170 QSOs per deg^2 from a sample of about 260 targets per deg^2 , for a target selection efficiency (including redshift failures) of 67%. For the subsequent passes, the target selection efficiency will be 100%, as only objects identified as $z > 2.1$ QSOs will be re-observed. After four passes, the average target selection efficiency is therefore of 88%.

3.4.4 Near- and mid-term developments for QSO target selection

The near-term focus (May-October 2015) is to build a large truth table of QSOs against which to test current and improved selection algorithms. BOSS and eBOSS surveys have both dedicated ancillary programs for comprehensive selections of quasars using the deep and multi-epoch SDSS photometry in the Southern Equatorial region called Stripe 82, where variability selections are notably efficient (cf. Sec. 3.4.5 and [217, 216]). The preliminary analysis of these data yields a sample of about 18,000 spectroscopically identified QSOs, over 120 deg^2 , to an extinction-corrected magnitude $g_c < 22.5$. These programs thus provide a sample with densities of 104 and 44 QSOs per deg^2 in the redshift ranges $z < 2.1$ and $z > 2.1$, respectively. Comparison to recent QSO luminosity functions [216] indicates that this sample is 85% complete. It thus constitutes a valuable truth table to probe QSO target selection to $g_c < 22.5$. This large sample selected almost solely from a well-controlled variability algorithm will also be used to improve the measure of the QSO luminosity function, in particular towards faint magnitudes. Before the end of 2015, we will extend this truth table with dedicated programs aiming at completing the selection of QSOs in $22 < g_c < 23$. Additional spectroscopic observations over 10 deg^2 should allow us to identify a total of about one thousand QSOs in this magnitude range.

In parallel, work has begun on selection algorithms such as XDQSO, to take better advantage of the imaging data available for DESI. In particular, we plan to use updated QSO luminosity functions to extend the range of validity of the code beyond $g_c = 22$, and to use the new truth tables to optimize the selection with grz +WISE photometry. Given the near 20% improvement observed in BOSS with the use of such algorithms, we can reasonably expect to increase the yield of the QSO selection for DESI by a similar factor. These developments will start with DES and DECaLS optical data, and existing WISE infrared data. They will be iterated as additional depth is acquired on WISE.

3.4.5 Variability Data Improves Selection of High-Redshift QSOs

Time-domain photometric measurements can enhance QSO selection. They allow us to exploit the intrinsic variability of QSOs [229] to distinguish them from stars of similar colors. They therefore complement the color-selection techniques presented in Sec. 3.4.2. We expect to use variability information in a first step to build truth tables against which to test QSO selection, and in a second step to select additional high-redshift Ly- α QSOs, for which uniformity of selection across the sky is not required.

Because the accretion region around a quasar is highly compact, its luminosity can vary substantially on timescales ranging from days to years, with a pattern distinct from that seen in variable stars. The time-variability of astronomical sources can be described using a measure of the amplitude of the observed variability as a function of the time delay between two observations. This “structure function” is modeled as a power law parameterized in

terms of A , the mean variation amplitude on a one-year time scale (in the observer’s reference frame) and γ , the logarithmic slope of the variation amplitude with respect to time.

We have tested variability techniques in DESI pilot surveys, both in Stripe 82 [217] that was the subject of repeated SDSS observations totaling about 50 epochs, and elsewhere on the sky, where time-domain information was derived from 5-10 epochs of PTF R -band data. As illustrated in Figure 3.18, the segregation between QSOs and stars is much reduced with poorer data, but variability remains competitive. This technique allowed us to identify 30% more QSOs in the Stripe 82 field than with time-averaged optical photometry only [217], and a combined color and variability selection from CFHT and PTF imaging data in the CFHTLS D3 field allowed us to achieve a record-high surface density of 207 QSOs per deg^2 to $g = 23$. The gain relative to the baseline QSO targeting with full *WISE* depth is likely to be less dramatic, and will be evaluated as those data become available.




Figure 3.18: Structure function parameters for 50-epoch gri light curves from SDSS in Stripe 82 (left), and for 6-epoch R light curves from PTF (right). The discriminating power is clearly reduced when fewer epochs and filters are available, but time-domain information remains valuable for identifying QSOs.

Imaging surveys that could provide useful time-domain information for variability selection include the PTF and follow-on iPTF surveys (in the deepest areas of their footprint), the future ZTF survey, the DES survey and DECam legacy survey, or a proposed wide-field imaging survey on CFHT. Variability is not assumed in our baseline targeting plan, but it is expected to be valuable for selecting the Ly- α QSOs wherever coverage exists.

3.5 Calibration Targets

Target selection is also responsible for providing lists of standard stars for flux calibration, and lists of blank sky locations to be used for modeling the sky.

Main-sequence F stars will be used as the primary spectrophotometric standard stars. These stars are well-described by stellar atmosphere models, making them ideal targets for spectrophotometric calibration at optical wavelengths. A stellar template of appropriate temperature, surface gravity and metallicity will be derived for each star and used to derive the spectral response including the time-varying atmospheric absorption bands.

The selection will be similar to the color-magnitude selection of BOSS to identify low-metallicity targets through a selection in $(u - g)$, $(g - r)$, $(r - i)$, and $(i - z)$ colors. The restrictive BOSS selection yields 10 stars per deg^2 ; to obtain a larger number of potential

targets using the new *grz* photometry, DESI will broaden this selection and include higher metallicity standard stars. With Gaia spectrophotometry of F stars that span a range of metallicity, and upcoming data from SDSS-IV/eBOSS in which a broader selection is applied, we plan to evaluate the value of a mix of lower and higher metallicity F stars to serve as flux calibration standards for DESI. Finally, we will perform a cross-calibration of low-metallicity and higher metallicity F-stars during the commissioning stages of DESI, thus providing validation of the standard star selection.

Blank sky locations will be determined as part of the object detection algorithms applied to the input imaging, ensuring that there are no detectable sources within the fiber diameter in any of the input bands. These will be provided at a density such that every fiber (when possible) will have the option of a blank sky if it isn't otherwise assigned to a science target.

3.6 Baseline Imaging Datasets

The samples described above can be selected given highly-uniform optical imaging data in the *g*, *r*, and *z* bands, as well as all-sky imaging from the *WISE* satellite. The same imaging data for selected science targets will be used to identify calibration targets (standard stars and sky fibers) as discussed in the accompanying TDR. A combination of three telescopes will be used to provide the baseline targeting data for DESI: the Blanco 4m telescope at Cerro Tololo, the Bok 90-inch and the Mayall 4m telescope at Kitt Peak. The footprints of the primary surveys using these telescopes that will deliver the targeting data are shown in Figure 3.19 and the next three subsections discuss these surveys and their current status in more detail. The status of the *WISE* data is presented in § 3.6.4.

3.6.1 Blanco/DECam Surveys ($\text{DEC} \leq 30^\circ$)

The Dark Energy Camera (DECam) on the Blanco 4m telescope, located at the Cerro Tololo Inter-American Observatory, will provide the optical imaging for targeting over 2/3 of the DESI footprint, covering both the North and South galactic cap regions at $\text{Dec} \leq 30^\circ$. Due to the combination of large field of view and high sensitivity from 400-1000 nm, DECam is the most efficient option for obtaining photometry in the *g*, *r*, and *z* bands.

DECam can reach the required depths for DESI targets in modest total exposure times of 100, 100 and 200 sec in *g*, *r*, *z* in median conditions. In median conditions, these data reach 6σ depths of $g=24.0$, $r=23.4$ and $z=22.5$ for an ELG galaxy with half-light radius of 0.45 arcsec. For a 3-dither observing strategy, accounting for weather loss, DECam is capable of imaging 9000 deg^2 of the DESI footprint to depth in 81 scheduled nights. These depth estimates have been vetted with *grz* photometry in the COSMOS field in Spring 2013 (Section 3.3.1).

A public survey, “The DECam Legacy Survey of the SDSS Equatorial Sky” (DESI collaborators D. Schlegel and A. Dey are PIs), has been approved to obtain optical imaging to the required depth over 6200 deg^2 . The survey (known as “DECaLS”) will be allocated 64 nights spread out over 3 years (2014A to 2017B semesters). The survey began in August 2014 and has thus far had 18.9 usable nights (out of the 20 assigned) during which 25% of the *g/r* and 35% of the *z* imaging was completed. The current coverage is shown in figure 3.20.

The DECaLS data resulting from the 2014B semester observations have been reduced: pipeline processing was done at NOAO and the calibrations and catalog construction (using the Tractor algorithm; see § 3.8) were performed at NERSC. Calibrated data and catalogs




Figure 3.19: The primary imaging surveys that will result in targeting data for the DESI project. The footprint at $\text{DEC} \leq +30^\circ$ will be covered using the Dark Energy Camera (DECam) on the Blanco 4m telescope at Cerro Tololo Inter-American Observatory. The Dark Energy Camera Legacy Survey (DECaLS, shown in yellow) and the Dark Energy Survey (DES, shown in orange) are underway; a proposal to cover the region labelled DECaLS+ (shown in purple) is pending. The footprint at $\text{DEC} \geq +30^\circ$ (in the North Galactic Cap region, shown in cyan) will be targeted using the 90Prime camera at the Bok 2.3-m telescope for the g and r band imaging (BASS: the Beijing-Arizona Sky Survey) and the MOSAIC camera on the Mayall 4m telescope for the z -band imaging (MzLS: the MOSAIC z -band Legacy Survey). Both the Bok and Mayall telescopes are located on Kitt Peak.

have been publicly released (see <http://legacysurvey.org>) and the DESI collaboration has begun tests of target selection on these data.

DECaLS will only cover $\approx 2/3$ of the planned DESI footprint at $\text{Dec} \leq 30^\circ$. An additional 500 deg^2 in the SGC will be observed to even greater depth with the same camera by the Dark Energy Survey; these data will be reprocessed (to ensure consistency with the rest of the footprint) by the DECaLS collaboration, and will be released publicly as part of the DECaLS data releases. The remaining $\approx 3000 \text{ deg}^2$ of the DESI footprint requires an additional 26-night allocation of time. We are in the process of exploring various avenues, including expanding the current DECaLS project to cover this area.




Figure 3.20: *Left panel:* Coverage map of the DECaLS survey based on data taken during the 2014B and 2015A semesters. The regions covered by 1, 2 and 3 passes are represented as open black, filled green and filled blue circles. 45 nights remain to be allocated, spread over the 2015B to 2017A semesters. *Right panel:* Coverage map of the Bok/BASS survey based on data collected over 56 nights during the 2015A semester. The regions covered by 1, 2 and ≥ 3 passes are represented as open black, filled green and filled red circles. 144 nights remain to be allocated.

3.6.2 Bok/90Prime Survey ($\text{DEC} \geq 30^\circ$)

The NGC footprint at $\text{Dec} \geq +30 \text{ deg}$ will be observed by the Bok 2.3-m telescope in two optical bands (g and r) for DESI targeting. The Bok Telescope, owned and operated by the University of Arizona, is located on Kitt Peak, adjacent to the Mayall Telescope. The 90Prime instrument is a prime focus $8\text{k} \times 8\text{k}$ CCD imager, with four University of Arizona ITL $4\text{k} \times 4\text{k}$ CCDs that have been thinned and UV optimized with peak QE of 95% at 4000\AA

and about 7 electron readout noise [230]. These CCDs were installed in 2009 and have been operating routinely since then. 90Prime delivers a 1.12 deg field of view, with 0.45'' pixels, and 94% filling factor. Typical delivered image quality at the telescope is 1.5''. The g and r -band survey over 5000 deg² requires 180 nights of scheduled telescope time. The throughput and performance in these bands were demonstrated with data in September 2013.

The Bok survey (known as the Beijing-Arizona Sky Survey; Zhou Xu and Xiaohui Fan, PIs; see <http://batc.bao.ac.cn/BASS>) has been awarded 200 nights (+40 for contingency) through Spring 2017. The Bok survey will target 5500 deg² in the NGC, including 500 deg² of overlap with the region covered by the DECam surveys in order to understand and correct for any systematic biases in the target selection. Through April 2015 the Bok survey has completed 42 nights of observing with about 60% efficiency (out of 56 scheduled in 2015A). All of the imaging through April has been in the g -band while awaiting the delivery of a new r -band filter to match the DECam characteristics, which was delivered by Asahi in late April. The area covered has been primarily at Dec $< 50^\circ$ (see Figure), following the same strategy as DECaLS of dividing the footprint into three passes, with at least one pass having photometric conditions (P1). Over half of the Bok area has been imaged with the g filter in P1 conditions, and just over a third of the total exposure requirement (including three passes) has been completed. Roughly three nights have been lost to issues that required a daytime fix, but no major issues have caused significant time loss. A number of instrument control software updates, new flexure maps, and new observing tools have been implemented that greatly improve the pointing accuracy, focusing of the telescope, and observing efficiency.

Analysis of data acquired during the January run shows that the survey is reaching the required g -band depth (see Figure 3.20). Pipeline development work is in progress and preliminary photometric catalogs have been produced, with the first data release planned for December 2015. The raw and calibrated data and catalogs will be publicly served through the NOAO Science Archive.

3.6.3 Mayall/MOSAIC Survey ((DEC $\geq 30^\circ$)

The z -band imaging survey of the northern galactic cap DESI footprint at DEC $\geq +30^\circ$ will be conducted using the MOSAIC3 camera at the prime focus of the 4-m Mayall telescope of the Kitt Peak National Observatory. The z -band survey has been assigned 230 nights during semesters 2016A and 2017A through an agreement between the National Science Foundation and the Department of Energy. NOAO has purchased a new z -band filter to match the DECam filter bandpass and to thereby minimize any differences between the DECam and MOSAIC z surveys. The KPNO 4m telescope control system, MOSAIC imaging camera software, and guiding and acquisition systems are being upgraded to run the survey as efficiently as possible.

The MOSAIC3 camera is a new version of the prime focus imaging system: the dewar is currently being refitted at Yale University with four 500 μ m-thick LBNL fully-depleted 4096² CCDs. The refitted camera will be delivered to NOAO in September 2015 and commissioned at the telescope in the 2015B semester. If, for any reason, the upgrade project is delayed, we will undertake the survey with the current version of the prime-focus camera (MOSAIC1.1); the required number of survey nights was determined using the current sensitivity of MOSAIC1.1, and the upgraded camera should result in a slightly deeper or wider survey. We anticipate the survey beginning in January 2016.

The MOSAIC z -band survey project will be run similarly to the DECaLS survey, with the initial processing being done using the NOAO pipeline and calibration and catalog construction being carried out at LBL/NERSC. The raw and pipeline-processed data will be public as soon as they are available, and catalogs will be released on a yearly basis. All data will be publicly available through the NOAO Science Archive and NERSC.

3.6.4 *WISE* All-Sky Survey

Infrared imaging from the Wide-field Infrared Survey Explorer (*WISE*) satellite are critical to the DESI targeting algorithm for LRGs and QSOs. During its primary 7-month mission from through August 2010, *WISE* conducted an all-sky survey in four bands centered at 3.4, 4.6, 12 and 22 μm (known as W1, W2, W3 and W4) [231] 99.99% of the sky was imaged at least 8 times, while regions near the ecliptic poles were observed more than 100 times. Following the primary 4-band mission, *WISE* continued survey operations in the three shortest bands for 2 months, then the two shortest bands for an additional 4 months. The 13 months of observations in the two shortest-wavelengths bands at 3.4 and 4.6 μm are used for DESI targeting. The limiting magnitudes (at 5σ) in the *WISE* All-Sky Data Release are 20.0 and 19.3 AB mag. DESI will use the re-stacked Level 1 *WISE* imaging that retain the intrinsic resolution of the data and are appropriate for photometry preserving the available signal-to-noise [232]. The photometry is measured using *the Tractor* as described in Section 3.8.

The *WISE* satellite will quadruple the exposure times in the W1 and W2 bands of interest for DESI. NASA reactivated the satellite on Fall 2013 and will continue two-band survey observations for an additional 3 years starting December 1, 2013, as the NEOWISE project. The first NEOWISE data release occurred in March 2015, with the final data release expected in 2017. These data will surpass the data used for DESI target selection depths by ~ 0.7 mag and will permit us to significantly improve the target selection beyond the baseline strategy.

3.7 Additional Imaging Data

Additional imaging data, if available, can supplement the target selection data and may be used, in particular, to improve the selection of the high-redshift Ly- α forest QSO sample. This is because the Ly- α forest analysis is based on the clustering of absorption systems along the line of sight, and therefore does not require a spatially uniform QSO sample. As a result, the QSO target selection can utilize datasets that may not be uniform (in depth, bandpass, or time sampling) over the DESI footprint. In this section, we summarize the key datasets that may contribute to this effort, if they prove to be available. These data sets are not assumed to be available for our baseline target selection plans, but rather should improve the efficiency of targeting higher-redshift ($z > 2.1$) QSOs beyond the baseline targeting strategy presented above.

Below we describe other wide field imaging surveys that are of interest for the DESI project, either for calibration purposes or as complementary data sources for high- z QSO selection.

3.7.1 SDSS

The Sloan Digital Sky Survey [233] has obtained multi-band (*ugriz*) photometry (in photometric conditions) over a 10,000 deg² extragalactic footprint in the North Galactic and South Galactic Caps. The Northern Cap and four stripes in the Southern Cap were imaged in 1998-2004. The bulk of the Southern Cap was imaged in 2008-2009, and the SDSS camera was then retired from service in December 2009. The median 5σ magnitude depths for the SDSS *ugriz* bands are 22.15, 23.13, 22.70, 22.20, and 20.71, respectively, but with substantial variation in depth from seeing. SDSS will provide a reference photometric point for variability selection of high-redshift QSOs, allowing variability over long time baselines to be measured.

3.7.2 PanSTARRS-1

The PanSTARRS-1 (PS1) 3π survey [234] is a transient-sensitive survey designed to observe 30,000 deg² of sky over 12 epochs in each of the five *grizy* survey filters. The multi-band photometry generated from the co-added exposures reaches depths that are comparable to SDSS in *gr* and potentially deeper in *iz*. This PS1 *z*-band photometry may be sufficiently deep for targeting DESI LRGs, but is insufficiently deep in the other bands for LRGs or in any bands for the DESI ELGs and QSOs. The PS1 survey completed observations in 2013. The co-added data and catalogs are expected to be publicly released by STScI/MAST in 2015. The PS1 time-domain photometry may be useful for enhancing the selection of Ly- α QSOs at the brighter magnitudes. The DECaLS survey is currently using the PS1 data to provide initial photometric and astrometric calibration across its footprint.

3.7.3 PTF, iPTF, and ZTF

The Palomar Transient Factory (PTF) [235] was a photometric survey designed to find transients via repeated imaging over 20,000 deg² in the Northern Hemisphere. In February 2013, the next phase of the program, iPTF (intermediate PTF) began. Both have used the CFH12K camera on the 1.2 m Oschin Telescope at Palomar Observatory, which covers 7.2 deg² of sky in a single pointing with a pixel scale of 1.01 arcsec.

Four years of survey operations have so far yielded a total of 5,000 deg² in *R*-band and 1,000 deg² in *g*-band to useful depths for QSO selection based on variability. LBNL is a partner in the PTF and iPTF collaborations, and DESI has access to these data.

The Zwicky Transient Factor (ZTF) is scheduled for first light in January 2017 with operations commencing within six months after being successfully funded through a Mid-Scale Innovations Program in Astronomical Sciences by NSF. They will use the same telescope with a new 46 square-degree imager. The ZTF survey will cover the entire sky at declinations $\text{Dec} > -20^\circ$, including the full DESI footprint. ZTF will operate with a *g*-band similar to the DECam and Bok *g*-band, an *R*-band (Mould-*R*) that is broader, and potentially an *i*-band. These data, which will be available to DESI collaboration for the purposes of target selection, are expected to eventually achieve the DESI targeting depths in *g* and *R* bands, but perhaps not before the start of DESI spectroscopic operations. The time sampling of ZTF is planned to be highly non-uniform over the DESI footprint, with different areas of sky covered in different years. Therefore, ZTF is not viable for the baseline DESI target selection, but PTF, iPTF and ZTF may be used to supplement the high-redshift QSO selection

for DESI. The independent coverage provided by ZTF can also be helpful in tying down the photometry across the entire DECaLS footprint.

3.7.4 CFHT

The Canada-France-Hawaii Telescope (CFHT) is a 3.6-m meter telescope on Mauna Kea, Hawaii. CFHT is a joint facility of the National Research Council of Canada, the Centre National de la Recherche Scientifique of France, and the University of Hawaii. The CFHT prime focus imager MegaCam, a very efficient instrument for imaging large areas of sky, consists of 36 $2k \times 4k$ e2v CCDs, covering a field of view of 0.97 deg^2 with a pixel scale of 0.185 arcsec per pixel. MegaCam started operations in 2003 and has conducted a number of large imaging surveys, the largest being the CFHT Legacy Survey covering 155 deg^2 .

The CFHT community is in discussions with the Euclid consortium and may play a role in providing *ugri* imaging data over the northern Euclid footprint. However, no plan is currently in place. There is an ongoing *u*-band survey (“CFHT-Luau: The CFHT Legacy Survey for the *u*-band all-sky universe”; A. McConnachie and R. Ibata, PIs) aimed at providing imaging over 4000 deg^2 of the high-Galactic-latitude northern sky, approximately split between the North and South Galactic caps. CFHT-Luau will complete in the 2016B semester (with data becoming public 1 year after observation). ($u - g$) color selection is an efficient discriminator between low-redshift and high-redshift QSOs. Hence, the CFHT data may be used to supplement the high-redshift Ly- α forest QSO selection in DESI, especially in combination with variability data.

3.7.5 SCUSS

The South Galactic Cap U-band Sky Survey [236] is a survey of 4000 deg^2 in the South Galactic Cap using the 90Prime instrument on the Bok 2.3-m telescope. The survey was a joint project among the Chinese Academy of Sciences, its National Astronomical Observatories unit, and Steward Observatory ⁶. The survey was conducted between September 2010 and October 2014 with typical exposure times of 5 minutes per field. The limiting magnitude reached by the data is $u \sim 23 \text{ mag}$ (5σ point source), with some variation due to varying seeing conditions. These data may be used to supplement the high-redshift Ly- α forest QSO selection in DESI, especially in combination with variability data.

3.8 The Tractor Photometry for Target Selection

The DESI target selection will combine photometry from the optical data and from *WISE*. DESI collaborator Dustin Lang has developed *the Tractor* forward-modeling approach to the pixel level data [237]. ⁷ This is a mathematically proper approach to fitting the differing PSF and pixel sampling of these data, where the optical data has a typical PSF of $\approx 1 \text{ arcsec}$ and the *WISE* PSF is $\approx 6 \text{ arcsec}$.

The Tractor takes as input the individual images from multiple exposures in multiple bands, with different seeing in each. A pixel-level model of the appearance of each source is fit to all the images simultaneously. Thus, if a source is determined to be a point source, it is photometered as a point source in every band and every exposure. If it is found to be a

⁶SCUSS survey <http://batc.bao.ac.cn/Uband/>

⁷<https://github.com/dstndstn/tractor>




Figure 3.21: An example “brick” covering $0.25 \times 0.25 \text{ deg}^2$ from the DECaLS survey. From left to right, the panels show the actual *grz* imaging data, the rendered model based on the Tractor catalog of the region, and the residual map. The Tractor catalog represents an inference-based model of the sky that best fits the observed data. All catalogs and data for the observed regions of sky are available at <http://legacysurvey.org>

galaxy with a given intrinsic light profile, that profile is consistently used in all bands. This produces object fluxes and colors that are as consistent as possible across these wide-area imaging surveys that necessarily have varying image quality.

For bright objects that were cleanly detected by *WISE* alone, we find our measurements to be consistent with the catalog measurements (see Figure 3.22). However, we are also able to measure the fluxes of significantly fainter objects, as well as to study collections of objects that are blended in the *WISE* imaging but resolved in the optical. As an example of how this increases the color-space information to which we have access, we show in Figure 3.23 a traditional “catalog-matching” optical-infrared color-magnitude diagram versus the results of our *WISE* forced photometry, which requires no matching.

The Tractor allows DESI target selection to utilize information from low signal-to-noise measurements. The differentiation between stars and QSOs relies upon *WISE* flux measures that can be below the 5σ detection limit of the official *WISE* catalogs (Sec. 3.4.2). *The Tractor* successfully differentiates between the QSOs that are detected in *WISE*, and the QSOs that in general are not detected (c.f., Figure 3.16). Traditional “catalog-matching” approaches would not be successful.

The Tractor has been used to produce photometry for DESI-like target selection of LRGs and QSOs as part of a BOSS ancillary program in Spring 2014. *The Tractor* has also been used for the primary target selection for LRGs and QSOs for the SDSS-IV/eBOSS that began observations in July 2014. For these targets, *the Tractor* forced photometry was applied based upon the SDSS pipeline-measured galaxy profiles. Those profiles were fixed, convolved with the *WISE* point-spread function, and then a linear fit was performed on the full set of *WISE* imaging data. The result is a set of flux estimates for all SDSS objects, constructed such that the sum of flux-weighted profiles best matches the *WISE* images. DESI will make use of these same fits, using DECam and Bok optical images rather than the SDSS images. Tractor catalogs based on the DR1 data release from the DECaLS project are already available publicly and are being used for tests of the baseline target selection process (see Figure 3.21).




Figure 3.22: Forced photometry results from *the Tractor* code, using information from SDSS detections and light profiles to measure the flux from objects in the *WISE* images to below the *WISE* detection limit. *Left:* The results agree for bright objects that are detected in the *WISE* catalog. The widening locus below $W1 \sim 14$ is due to our photometry treating larger objects as truly extended, in contrast to the point-source-only assumptions in the public *WISE* catalog. *Right:* A demonstration of the increased depth made possible from using *the Tractor*. By using optical imaging from SDSS to detect objects, photometry is measured for objects that are well below the *WISE* detection limit.




Figure 3.23: Forced photometry results from *the Tractor* code, contrasted with traditional “catalog-matching”. *Left:* Color-magnitude diagram from matching SDSS to *WISE* catalogs. Many objects below the *WISE* catalog detection limits are lost. *Right:* Results from forced photometry of the *WISE* images based on SDSS detections. No matching is required, and objects that would be detected in *WISE* at only few-sigma significance can readily provide flux measurements.

4 Survey Design

4.1 Introduction

The DESI instrument will make largest spectroscopic survey to date. The design of the survey is optimized by selecting a footprint that is as large as possible from the Mayall telescope while staying clear of the Milky Way. The survey strategy will establish the order in which the observations will be made. The strategy will be modified in detail by atmospheric conditions, but the overall plan will be established to optimize the best science results for both the complete survey and results from intermediate years.

4.2 Survey Footprint

The DESI survey footprint is defined to be 14,000 square degrees that can be observed spectroscopically from Kitt Peak. This footprint will be one contiguous region selected from the North Galactic Cap (NGC) and one contiguous region in the South Galactic Cap (SGC). The instrumented area of the focal plane is 7.50 square degrees. About 14,000 square degrees can be covered nearly completely with one pass of 2,000 tiles, where each tile represents one DESI observation. Five passes with altogether 10,000 tiles covers each coordinate of the footprint with an average of 5.24 fibers. The DESI footprint is formally defined as any position on the sky within 1.605 deg of any of these selected tile centers.

The DESI spectroscopic survey will primarily select targets from catalogs derived from imaging with the Blanco/DECam camera, the Bok/90Prime camera, and the Wide-field Infrared Survey Explorer (*WISE*). Although WISE imaging covers the entire sky, the imaging from DECam and the Bok Telescope impose an external constraint on the DESI footprint, as targets must be selected from large contiguous regions imaged with the same instruments. The Bok will provide targeting in the NGC at $\text{Dec} > +30$ deg. The Blanco will provide targeting in both the NGC and SGC at $\text{Dec} < +30$ deg. An area of approximately 800 sq. deg. in the SGC at $\text{Dec} > +30$ (and $-32 < b < -15$) is “orphaned” and excluded from the DESI survey as it would be a small area observed with a different camera.

The footprint is constrained, as well, by the need to avoid regions that would require long exposures due to airmass or dust, by weather patterns at Kitt Peak, and by regions of high stellar density. The resulting footprint is shown in Fig. 4.1.

4.3 Field Centers

We refer to “tiling” as the process by which field centers are assigned in a manner to cover the footprint with optimal coverage of each coordinate on the sky. The one-pass tiling of the sky mentioned in Section 4.2 is a preliminary solution that is achieved using the icosahedral tiling [238] with 5762 tile centers distributed on the full sphere⁸. This tiling is very-well matched to the DESI focal plane size. The first pass rotates the above tiling solution by 90 deg in RA. This rotation conveniently puts rows of tile centers along lines of approximately constant declination at the north and south boundaries of the DESI survey. Each of additional passes 2 through 5 have an additional rotation of the tile centers by 1.08 deg in RA. This gives large dithers on most of the sky (except at the pole, which is not in the DESI footprint), thus filling the gaps in the focal plane with subsequent visits. Non-uniformity in coverage

⁸<http://neilsloane.com/icosahedral.codes/>

could artificially introduce structure in the targeting of LSS-tracers; alternative tilings based on the same first pass but with subsequent passes obtained with more disparate rotations will be further studied for possible improvements to the uniformity.




Figure 4.1: Tile centers for the DESI footprint in an equal-area projection. Declination limits are imposed at $-8.2 < \text{Dec}$ in the NGC (left), and $-18.4 < \text{Dec} < +30$ in the SGC (right). Approximately 1% of tiles have exposure factors larger than 2.5 (shown in blue), but are included to avoid unwanted holes in the footprint. The five passes are shown in separate colors. The spots indicate the centers of focal plane positions, not the size of the area in the sky subtended by the focal plane. Locations inside the footprint are within reach of a fiber, on average, 5.24 times during the survey.

A different tiling solution is necessary for the survey with the reduced focal plane. With only six of the 10 wedges populated on the focal plane there are 3000 instead of 5000 fibers. The populated wedges are best arranged in a “Pacman” format. First, 240 tile centers are placed on the celestial equator uniformly separated in RA. Stripes of tiles are then placed on lines of constant celestial latitude spaced every 2.765 deg. At each stripe, the number of tiles is reduced by the factor $\cos(\text{Dec})$ from the 240 placed on the celestial equator. This results in a tiling solution with similar uniformity and coverage statistics as the baseline survey, with 4% more tiles than would be necessary under the assumption of a simple scaling with focal plane area.

The pattern of fiber positioners in the focal plane is shown in Fig. 4.2. Combining this with the tiling gives a purely geometric measure of the coverage for each position within the DESI footprint. The distribution of this coverage is shown in Fig. 4.3 and in Table 4.1. The average coverage is about 5.1, with only 3.5% of the footprint having a coverage of less than 3. The edges of the footprint have the least coverage.




Figure 4.2: Fiber positioner locations for the full DESI instrument (left) and the reduced instrument “pacman” configuration (right). “Missing” positioner locations are for the guide-focus arrays (square regions) and fiducial markers for the fiber view camera.

The results of a similar study for the reduced “Pacman” focal plane are shown in the right hand panel of Fig.4.3 and Table 4.1

4.4 Observation Strategy

4.4.1 Sequence of Observations

The placement of field centers presented in Section 4.3 is designed to cover the footprint in five independent tilings. Given the 1940 hours of scheduled time each year, roughly 20% (390 hours) will occur under grey conditions and the remainder under dark conditions. Each year, 20% of the fields in a full independent tiling of 2000 field centers will be observed using the scheduled time with the highest level of background from the moon. This pass is planned to include only ELG targets because their spectral features are predominantly found at redder wavelengths and redshift success rates are less susceptible to increased sky background from the moon. On the other hand, the darkest 80% of the scheduled time (1550 hours) will be used to observe the QSO and LRG targets at highest priority, leaving the remaining science fibers for ELG targets as described in Section 4.3.

There remains additional freedom to determine the order in which the tiles over the four dark time passes are observed. Full simulations of the program will be used to determine

Table 4.1: The fraction of the footprint covered by 1, 2, 3, ... 8 fibers after five passes with the baseline survey and after just the first four passes, the only ones observing LRGs and QSOs. The mean is slightly decreased and the rms slightly increased by edge effects.

Coverage	1	2	3	4	5	6	7	8	Mean	RMS
All five passes	0.016	0.019	0.040	0.155	0.424	0.279	0.055	0.009	5.06	1.175
Four passes only	0.021	0.037	0.152	0.482	0.462	0.040	0.004	0.000	4.04	1.012




Figure 4.3: Coverage pattern on the sky after five passes for a 4 degree by 4 degree patch. This is shown for a region away from the edges of the footprint. **Left:** the fully-populated focal plane with 5000 fibers. **Right:** the reduced focal plane with 3000 fibers.

the optimal approach. The simulations will factor in seeing, transparency and weather variations for each exposure via Monte Carlo simulations to predict the quality of spectra and the variations in final survey areal coverage. Each exposure will be tuned to a grid of targets parameterized by magnitude and redshift using an exposure time calculator that approximates the sensitivity of the instrument. Weather conditions will be mocked using monthly statistics at Kitt Peak and the results will be used to determine likely redshift success rates over all target classes. The approach that optimizes intermediate and final cosmology results will be chosen.

We provide two examples of strategies, evaluated without consideration of weather and other variables. The first strategy simply follows the pattern given by the five layers of the footprint, observing one layer during grey time and the other four layers sequentially. The second possible strategy is designed to get complete coverage of all target classes over one-tenth of the sky as quickly as possible. In both cases we establish the order of observations by going from lowest declination upwards.

The sequential survey is described by Table 4.2 and displayed in Fig. 4.4.

Table 4.2: The first survey strategy in which tiling layers 1–4 are observed sequentially. Pass 5 is observed in grey time and devoted to ELGs alone

	year1	year 2	year 3	year 4	year 5	total
Pass 1	1711	429	0	0	0	2139
Pass 2	0	1277	851	0	0	2128
Pass 3	0	0	854	1282	0	2136
Pass 4	0	0	0	426	1704	2130
Pass 5	426	426	426	426	426	2129

A survey designed to get an early complete sample over 10% of the footprint is described by Table 4.3 and shown in Fig. 4.5. As expected, after the first year (about 2000 tiles




Figure 4.4: The evolution of the observation of each galaxy time as a function of the number of tiles observed using the original five passes observed consecutively, except the fifth pass is done in grey time and some reordering is done to provide free fibers throughout the survey.

observed), the second scheme produces multiple observations of LRGs and QSOs that do not exist in the first strategy. The second strategy also produces complete coverage of QSOs, LRGs, and ELGs by the end of year two, whereas the first strategy does not obtain complete coverage over any area until year four. By the end of five years, both strategies should complete the same area with the same statistics on each target sample, assuming weather is as expected.

Table 4.3: The second survey strategy in which complete coverage of a 1400 sq deg footprint is prioritized. This is implemented by always working from lowest declination upward in each pass.

	year1	year 2	year 3	year 4	year 5	total
Pass 1	1283	856	0	0	0	2139
Pass 2	383	468	1277	0	0	2128
Pass 3	21	192	214	854	854	2136
Pass 4	21	192	213	852	852	2130
Pass 5	426	426	426	426	426	2129

4.4.2 Exposure Times and Margin

Over five years, DESI is projected to observe 14,000 sq. deg. of the footprint presented in Section 4.2. The exact subset of this footprint to be observed will be contiguous regions in each of the NGC and the SGC that best fit the expected allocation of time. We have simulated the choice of final tile centers and the average exposure times according to an observing schedule of 1940 hours of dark and grey time per year as defined in the Site Alternatives study (DESI-311). The simulation includes a two minute overhead between fields




Figure 4.5: The evolution of the observation of each galaxy time as a function of the number of tiles observed using the alternative scheme, which covers 10% of the footprint multiple times, completely, in the first two years. The fifth pass is done in grey time and some reordering is done to provide free fibers throughout the survey.

and variations in exposure time for each field due to airmass and Galactic dust extinction. All exposure times are split into two separate exposures (with one minute of read time). This split limits the number of cosmic rays in an individual exposure, and also effectively maximize the S/N in variable sky conditions. The accumulated S/N will be measured by the exposure time calculator (see the DESI Performance Studies in the Instrument TDR). We project that 57% of the scheduled time will deliver usable data, where “usable data” is assumed in conditions when the dome is open and seeing is better than 1.5 arcsec. Although DESI will observe when the seeing is worse than 1.5 arcsec, those data have been ignored in these estimates of survey duration.

We simulate the full suite of observations accounting for airmass and Galactic dust extinction by choosing an hour angle for each field that maximizes the overall survey depth while fitting into the allocated time. Exposure times are estimated for each field to produce uniform depth in dust-extinction and atmosphere-extinction corrected spectra. In preliminary estimates, we assume the same dependence of S/N on airmass as was measured with BOSS, and degradation in S/N due to Galactic extinction for the sky-noise-limited case of the faintest targets. In future iterations, we will include a more sophisticated interpretation of redshift success rate for representative targets, thus accounting for the wavelength-dependent S/N estimates of each target class. For the 14,000 sq. deg. footprint observed with 10,000 tiles, we find an average exposure time of 1800 seconds. Scaling this to an observation taken at zenith with no Galactic dust extinction (as shown in the Figure in the DESI Performance Studies in the Instrument TDR) produces an equivalent exposure time of 1226 seconds. In other words, each exposure will have a S/N equivalent to a 1226 second exposure taken at zenith, under photometric conditions, median sky brightness and median seeing. As explained in Simulations Section of the Instrument TDR this fiducial exposure time of 1226

seconds allows the 1000 second exposures that are predicted to produce the required redshift success rates for each DESI target class. This projection leaves a 22% margin in exposure time for worse-than-projected weather, throughput performance, instrument downtime, or other factors that could slow the pace of the survey.

Similarly, we have estimated the average and fiducial exposure times for the reduced focal plane of the DESI KPP survey. The “Pacman” tiling of Section 4.2 leads to an average exposure time of 1700 seconds for 10,600 tile centers covering 9,000 sq. deg. Even though the average exposure time is somewhat lower than the 14,000 sq. deg. survey, the fiducial exposure time of 1270 seconds is actually larger because the average field in a 9,000 sq. deg. program lies at lower airmass and lower Galactic extinction than the average field in a 14,000 sq. deg. program. The projected margin for the 9,000 sq. deg. KPP survey is 27%.

4.5 The Bright Galaxy and Milky Way Surveys

4.5.1 Introduction

A portion of DESI operations will be affected by increased sky brightness from the moon, so as to make conditions unsuitable for observing the targets above $z = 0.6$. DESI expects to observe in the darkest 21 nights of the month, but some of those nights are affected in part by moon, adding up to about 440 hours per year of time. Assuming the same average weather statistics used in planning for dark time, we expect 250 hours per year on average of open-dome bright time. During this time, the DESI collaboration will conduct a survey of bright galaxies which will increase performance for the cosmology goals. This Bright Galaxy Survey (BGS) will be the primary bright-time survey program. In addition, the density of fibers in the DESI focal plane will enable a simultaneous survey of Milky Way Stars (The Milky Way Survey; MWS) during bright time. The MWS will target some of the oldest stars in the Galaxy with the goal of understanding the mass distribution, formation and evolution of the Galaxy. We refer to these two combined programs as the Bright Time Survey (BTS).

4.5.2 Survey Footprint

The Bright Time Survey will use the same 14,000 square degree footprint as the dark time project. This will enable the BTS to benefit from the optimization of the dark time footprint for observability. The BGS targets will be selected from the same imaging data as the dark time targets. The MWS will use Gaia photometry and proper motions for target selection. The Gaia survey is all-sky, and so covers the DESI 14,000 square degree footprint.

4.5.3 Field Centers

The BTS will use the same tiling pattern as the DESI Key Project, but with only 3 layers totalling 6000 tiles. There are roughly 800 galaxies per square degree to $r = 19.5$. With 4500 science fibers per tile, the BTS will place about 27 million fibers, far more than the ~ 10 million BGS targets. However, the clustering and Poisson fluctuations of bright galaxies means that we must incur these extra layers if we want to achieve a higher completeness. Preliminary simulations of DESI fiber assignments for $r < 19.5$ galaxies from the SDSS using the DESI tiling show that two passes gives 90% completeness and three passes over the footprint will give 97–98% completeness.

There are about 600 stars with effective temperatures warmer than 4700 K per square degree to $r = 18$ at Galactic latitude greater than 40 degrees from the equator. The DESI focal plane is 7.5 square degrees, so at each pass there will be many fibers available for the MWS.

4.5.4 Observation Strategy

Completing 6000 tiles in the 1250 hours of available open-dome time indicates an average time of 12.5 minutes per tile. Survey simulations accounting for the increased exposure time required as a function of airmass and extinction indicate that we would have 400 seconds available for a reference exposure at unit airmass and zero extinction. We are planning for a 300 second reference exposure, therefore leaving a 33% margin. Our spectral simulations (§3.1 and Fig. 3.3) indicate that a 5-minute exposure will yield a redshift success of 95% for galaxies down to $r = 19.5$.

For stars, 5 minute exposures will result in spectra with S/N per Angstrom of 14 at $\lambda > 650$ nm for stars of r magnitude 16.5-18, depending on their spectral energy distributions. Spectra of that quality are sufficient to yield radial velocity and chemical abundance information.

Because the BTS needs only three passes, it will be possible to combine multiple exposures for fainter objects for higher S/N. For example, it would be possible to re-expose many of the 5% of BGS targets that fail to achieve a redshift in the first two layers. We will perform fiber assignment simulations that combine the MWS and BGS samples to determine the optimal way to assign fibers that accounts for galaxy clustering and the variation in stellar density across the footprint, and which achieves maximum redshift and radial velocity completeness for faint targets.

With this basic strategy we expect to obtain spectra of roughly 10 million galaxies in the BGS and 10 million stars in the MWS. More simulations of the BTS are required to determine how to prioritize sky coverage versus completeness to enable early science. The BTS simulations will use the same survey simulation code as the dark time program, adapting it as required.

A Acronyms

1D	One-Dimensional
3D	Three-Dimensional
A&G	Acquisition and Guiding
A/R	AntiReflection
AAO	Australian Astronomical Observatory
AAT	Anglo-Australian Telescope
ABL	AirBorne Laser Aircraft
ACT	Atacama Cosmology Telescope
ADC	Atmospheric Dispersion Compensator
ADC	Analog to Digital Converter
ADR	Atmospheric Differential Refraction
AGN	Active Galactic Nucleus
AI&T	Assembly, Integration and Test
AOS	Active Optics System
AP	Alcott-Pachinko [sic]
API	Application Programming Interface
APO	Apache Point Observatory
APOGEE	The APO Galactic Evolution Experiment
AR	Antireflection
ASTM	American Society for Testing and Materials
AURA	Association of Universities for Research in Astronomy
BAO	Baryonic Acoustic Oscillations
BCAM	Brandeis CCD Angle Monitor
BOSS	Baryon Oscillation Spectroscopic Survey
BTGS	Bright Time Galaxy Survey
CABT	Centrale d'Aquisition Basses Température (Low Temperature Data Controller)
CAD	Computer-Aided Design
CAN	Controller Area Network
CCD	Charge Coupled Device
CDE	Cooler Drive Electronics
CDM	Cold Dark Matter
CDR	Critical Design Review; Conceptual Design Report
CEA	Commissariat à l'Énergie Atomique
CFHT	Canada-France-Hawaii Telescope
CFHTLS	CFHT Legacy Survey
CM	Configuration Management
CMASS	BOSS Constant Mass Sample
CMB	Cosmic Microwave Background
CMM	Coordinate Measurement Machine
CNC	Computer Numerical Control
CoDR	Concept Design Review/Report
CPU	Central Processing Unit

CSS	Cascading Style Sheets
CTE	Coefficient of Thermal Expansion
CTE	Charge Transfer Efficiency (CCD)
CTIO	Cerro Tololo Inter-American Observatory
DAQ	Data AcQuisition
DE	Dark Energy
DEC	Declination
DECam	Dark Energy Camera
DEEP2	Deep Extragalactic Evolutionary Probe 2
DEIMOS	DEep Imaging Multi-Object Spectrograph
DES	Dark Energy Survey
DESI	Dark Energy Spectroscopic Instrument
DETF	Dark Energy Task Force
DGP	DvaliGabadadzePorrati
DIMM	Differential Image Motion Monitor
DOE	Department of Energy
DOF	Degrees of Freedom
DRP	Data reduction pipeline
DR9	SDSS-III/BOSS Data Release 9
DR11	SDSS-III/BOSS Data Release 11
DTS	Data Transport System
EE50	50% Encircled Energy
EE95	95% Encircled Energy
EE99	99% Encircled Energy
EGSE	Electrical Ground Support Equipment
ELG	Emission Line Galaxy
EPA	Environmental Protection Agency
ESNET	Energy Sciences Network
FEA	Finite Element Analysis
FEE	Front-End Electronics
FITS	Flexible Image Transport System
FLA	Field Lens Assembly
FNAL	Fermi National Accelerator Laboratory
FoM	Figure of Merit
FOV	Field of View
FPA	Focal Plane Assembly
FPD	Focal Plate Adapter
FPGA	Field Programmable Gate Array
FPS	Focal Plane System
FRD	Focal Ratio Degradation
FTS	Fourier Transform Spectrograph
FVC	Fiber View Camera
FWHM	Full Width at Half Maximum
GAMA	Galaxy And Mass Assembly Survey
GFA	Guide, Focus, and Alignment
GNU	GNU's Not Unix!

GUI	Graphical User Interface
HPSS	High Performance Storage Systems
HTML	HyperText Markup Language
HVAC	Heating, Ventilation, and Air Conditioning
I&T	Integration and Test
IAA	Instituto de Astrofísica de Andalucía
IAC	Instituto de Astrofísica de Canarias
ICD	Interface Control Document
ICS	Instrument Control System
IDD	Interface Definition Drawing/Document
IDL	Interactive Data Language
IGM	Intergalactic Medium
IP	Internet Protocol
IPAC	Infrared Processing and Analysis Center
iPTF	Intermediate Palomar Transient Factory
IR	InfraRed
KPNO	Kitt Peak National Observatory
L1	Level 1 requirements
L2	Level 2 requirements
L3	Level 3 requirements
LAM	Laboratoire d'Astrophysique de Marseille
LBNL	Lawrence Berkeley National Laboratory
LBT	Large Binocular Telescope
LED	Light Emitting Diode
LHC	Large Hadron Collider
LOWZ	BOSS Low Redshift Galaxy Sample
LPT	Linear Pulse Tube
LRG	Luminous Red Galaxy
LSS	Large Scale Structure
LSST	Large Synoptic Survey Telescope
M2	Existing secondary mirror at Mayall 4m telescope
MAST	Mikulski Archive for Space Telescopes
MB	Megabyte
MDLF	Mikulski Archive for Space Telescopes
MG	Modified Gravity
MMT	Multiple Mirror Telescope
MOS	Multi-Object Spectrograph
MOU	Memorandum of Understanding
MPC	Multi-Point Constraint
MTBF	Mean Time Between Failures
MVC	Model-View-Controller
NA	Not Applicable
NASA	National Aeronautics and Space Administration
NCSA	National Center for Supercomputing Applications
NERSC	National Energy Research Scientific Computing Center
NFPA	National Fire Protection Association

NGC	Northern Galactic Cap
NIR	Near-Infrared
NOAO	National Optical Astronomy Observatory
NOAONVO	NOAO National Virtual Observatory
NOMAD	Naval Observatory Merged Astrometric Dataset
NSF	National Science Foundation
NVO	National Virtual Observatory
OCS	Observation Control System
ODI	One Degree Imager
OPD	Optical Path Difference
OPO	Optical Parametric Oscillator
OSHA	Occupational Safety and Health Administration
OSS	Observatory System Specification
PAH	Polycyclic Aromatic Hydrocarbon
Pan-STARRS	The Panoramic Survey Telescope and Rapid Response System
PC	Personal Computer
PCA	Principal Component Analysis
PDR	Preliminary Design Review
PF	Prime Focus
PFA	Positioner Fiber Assembly
PFC	Prime Focus Corrector
PFI	Prime Focus Instrument
PFU	Prime Focus Unit
PI	Principal Investigator
PLC	Programmable Logic Controller
PM	Project Manager/Management
PML	Physical Markup Language
POS	Fiber Positioner
PPM	Parts Per Million
PSF	Point Spread Function
PSM	Point Source Microscope
PTF	Palomar Transient Factory
PTL	Focal Plate Petal
PVC	PolyVinyl Chloride
PYRO	Python Remote Objects
QA	Quality Assurance
QC	Quality Control
QE	Quantum Efficiency
QSO	Quasi-Stellar Object
R&D	Research and Development
RA	Right Ascension
RFP	Request for Proposal
RGB	Red Giant Branch
RMI	Remote Method Invocation
RMS	Root Mean Square
ROC	Radius of Curvature

ROI	Region of Interest
ROM	Rough Order of Magnitude
RSD	Redshift Space Distortions
RSS	Root Sum Squared
RTD	Resistance Temperature Detector
RTV	Room Temperature Vulcanizing silicone
S/N	Signal to Noise
SCUSS	The South Galactic Cap U-band Sky Survey
SDF	Spectral Density Function
SDSS	Sloan Digital Sky Survey
SE	Systems Engineering
SED	Spectral Energy Distributions
SEGUE	The Sloan Extension for Galactic Understanding and Exploration
SEP	Systems Engineering Plan
SEQUELS	Sloan Extended QUasar, ELG and LRG Survey
SET	Systems Engineering Team
SGC	Southern Galactic Cap
SHE	Safety, Health and Environmental
SIDE	Super Ifu Deployable Experiment
SMBH	SuperMassive Black Hole
SN	SuperNova
SNR	Signal-to-Noise Ratio
SOW	Statement of Work
SPT	South Pole Telescope
SQL	Structured Query Language
SRD	Science Requirements Document
SRR	System Requirements Review
SRS	Software Requirements Specification
SVE	Shared Variable Engine
TB	TerraBytes
TBA	To Be Announced
TBC	To Be Confirmed
TBD	To Be Defined/Decided/Determined
TBR	To Be Reviewed/Revised/Resolved
TCP	Transmission Control Protocol
TCS	Telescope Control System
TES	Top-End Structure
TKRS	The Team Keck Treasury Redshift Survey
TTL	Transistor-Transistor Logic
UCL	University College London
UK	United Kingdom
UPS	Uninterruptible Power Supply
URL	Uniform Resource Locator
USA	United States of America
USB	Universal Serial Bus
USTC	University of Science and Technology China

UV	Ultraviolet
UVES	Ultraviolet and Visual Echelle Spectrograph
VAC	Volts AC
VIMOS	VIisible MultiObject Spectrograph
VIPERS	VIMOS Public Extragalactic Redshift Survey
VISTA	Visible and Infrared Survey Telescope for Astronomy
VLT	Very Large Telescope
VPH	Volume Phase Holographic
VVDS	VIMOS VLT Deep Survey
WBS	Work Breakdown Structure
WFC	Wide Field Corrector
WFIRST	The Wide-Field Infrared Survey Telescope
WFMOS	Wide-Field Multi-Object Spectrograph
WGL	Weak Gravitational Lensing
WIRO	University of Wyoming's 2.3 m telescope
WISE	Wide-field Infrared Survey Explorer
WIYN	Wisconsin, Indiana, Yale, NOAO Telescope
WMAP	Wilkinson Microwave Anisotropy Probe
WP	Wilkinson Microwave Anisotropy Probe
ZTF	Zwicky Transient Facility

References

- [1] S. Perlmutter et al. “Measurements of Omega and Lambda from 42 High-Redshift Supernovae”. In: *Astrophys. J.* 517 (June 1999), pp. 565–586. eprint: [astro-ph/9812133](#).
- [2] A. G. Riess et al. “Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant”. In: *Astron. J.* 116 (Sept. 1998), pp. 1009–1038. eprint: [astro-ph/9805201](#).
- [3] M. Levi et al. “The DESI Experiment, a whitepaper for Snowmass 2013”. In: *ArXiv e-prints* (Aug. 2013). arXiv: [1308.0847 \[astro-ph.CO\]](#).
- [4] A. Albrecht et al. “Report of the Dark Energy Task Force”. In: *ArXiv Astrophysics e-prints* (Sept. 2006). eprint: [astro-ph/0609591](#).
- [5] Daniel J. Eisenstein and Wayne Hu. “Baryonic features in the matter transfer function”. In: *Astrophys. J.* 496 (1998), p. 605. arXiv: [astro-ph/9709112 \[astro-ph\]](#).
- [6] L. Anderson et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: baryon acoustic oscillations in the Data Releases 10 and 11 Galaxy samples”. In: *Mon. Not. R. Astron. Soc.* 441 (June 2014), pp. 24–62. arXiv: [1312.4877](#).
- [7] F. Beutler et al. “The 6dF Galaxy Survey: baryon acoustic oscillations and the local Hubble constant”. In: *Mon. Not. R. Astron. Soc.* 416 (Oct. 2011), pp. 3017–3032. arXiv: [1106.3366 \[astro-ph.CO\]](#).
- [8] C. Blake et al. “The WiggleZ Dark Energy Survey: joint measurements of the expansion and growth history at $z \leq 1$ ”. In: *Mon. Not. R. Astron. Soc.* 425 (Sept. 2012), pp. 405–414. arXiv: [1204.3674 \[astro-ph.CO\]](#).
- [9] D. J. Eisenstein et al. “Improving Cosmological Distance Measurements by Reconstruction of the Baryon Acoustic Peak”. In: *Astrophys. J.* 664 (Aug. 2007), pp. 675–679. eprint: [arXiv: astro-ph/0604362](#).
- [10] N. Padmanabhan et al. “A 2 per cent distance to $z = 0.35$ by reconstructing baryon acoustic oscillations - I. Methods and application to the Sloan Digital Sky Survey”. In: *Mon. Not. R. Astron. Soc.* 427 (Dec. 2012), pp. 2132–2145. arXiv: [1202.0090 \[astro-ph.CO\]](#).
- [11] A. J. Ross et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: analysis of potential systematics”. In: *Mon. Not. R. Astron. Soc.* 424 (July 2012), pp. 564–590. arXiv: [1203.6499 \[astro-ph.CO\]](#).
- [12] B. Leistedt and H. V. Peiris. “Exploiting the full potential of photometric quasar surveys: optimal power spectra through blind mitigation of systematics”. In: *Mon. Not. R. Astron. Soc.* 444 (Oct. 2014), pp. 2–14. arXiv: [1404.6530](#).
- [13] Planck Collaboration et al. “Planck 2013 results. XVI. Cosmological parameters”. In: *ArXiv e-prints* (Mar. 2013). arXiv: [1303.5076 \[astro-ph.CO\]](#).
- [14] D. Eisenstein and M. White. “Theoretical uncertainty in baryon oscillations”. In: *Phys. Rev. D* 70.10, 103523 (Nov. 2004), p. 103523. eprint: [arXiv:astro-ph/0407539](#).
- [15] H.-J. Seo et al. “Nonlinear Structure Formation and the Acoustic Scale”. In: *Astrophys. J.* 686 (Oct. 2008), pp. 13–24. arXiv: [0805.0117](#).
- [16] M. Crocce and R. Scoccimarro. “Nonlinear evolution of baryon acoustic oscillations”. In: *Phys. Rev. D* 77.2 (Jan. 2008), pp. 023533–+. eprint: [arXiv:0704.2783](#).

- [17] R. E. Smith, R. Scoccimarro, and R. K. Sheth. “Motion of the acoustic peak in the correlation function”. In: *Phys. Rev. D* 77.4 (Feb. 2008), pp. 043525–+. eprint: [arXiv:astro-ph/0703620](#).
- [18] N. Padmanabhan and M. White. “Calibrating the baryon oscillation ruler for matter and halos”. In: *Phys. Rev. D* 80.6, 063508 (Sept. 2009), p. 063508. arXiv: 0906.1198 [[astro-ph.CO](#)].
- [19] H.-J. Seo et al. “High-precision Predictions for the Acoustic Scale in the Nonlinear Regime”. In: *Astrophys. J.* 720 (Sept. 2010), pp. 1650–1667. arXiv: 0910.5005 [[astro-ph.CO](#)].
- [20] K. T. Mehta et al. “Galaxy Bias and Its Effects on the Baryon Acoustic Oscillation Measurements”. In: *Astrophys. J.* 734, 94 (June 2011), p. 94. arXiv: 1104.1178 [[astro-ph.CO](#)].
- [21] B. D. Sherwin and M. Zaldarriaga. “Shift of the baryon acoustic oscillation scale: A simple physical picture”. In: *Phys. Rev. D* 85.10, 103523 (May 2012), p. 103523. arXiv: 1202.3998 [[astro-ph.CO](#)].
- [22] D. Tseliakhovich and C. Hirata. “Relative velocity of dark matter and baryonic fluids and the formation of the first structures”. In: *Phys. Rev. D* 82.8, 083520 (Oct. 2010), p. 083520. arXiv: 1005.2416 [[astro-ph.CO](#)].
- [23] J. Yoo, N. Dalal, and U. Seljak. “Supersonic relative velocity effect on the baryonic acoustic oscillation measurements”. In: *J. Cosmology Astropart. Phys.* 7, 018 (July 2011), p. 18. arXiv: 1105.3732 [[astro-ph.CO](#)].
- [24] L. Anderson et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations in the Data Release 9 Spectroscopic Galaxy Sample”. In: *ArXiv e-prints* (Mar. 2012). arXiv: 1203.6594 [[astro-ph.CO](#)].
- [25] C. L. Reichardt et al. “New Limits on Early Dark Energy from the South Pole Telescope”. In: *Astrophys. J. Lett.* 749, L9 (Apr. 2012), p. L9. arXiv: 1110.5328 [[astro-ph.CO](#)].
- [26] U. Alam. “Constraining Perturbative Early Dark Energy with Current Observations”. In: *Astrophys. J.* 714 (May 2010), pp. 1460–1469. arXiv: 1003.1259 [[astro-ph.CO](#)].
- [27] J.-Q. Xia and M. Viel. “Early dark energy at high redshifts: status and perspectives”. In: *J. Cosmology Astropart. Phys.* 4, 002 (Apr. 2009), p. 2. arXiv: 0901.0605 [[astro-ph.CO](#)].
- [28] E. V. Linder and G. Robbers. “Shifting the Universe: early dark energy and standard rulers”. In: *J. Cosmology Astropart. Phys.* 6, 004 (June 2008), p. 4. arXiv: 0803.2877.
- [29] M. Doran and G. Robbers. “Early dark energy cosmologies”. In: *J. Cosmology Astropart. Phys.* 6, 026 (June 2006), p. 26. eprint: [arXiv:astro-ph/0601544](#).
- [30] M. Doran, J.-M. Schwindt, and C. Wetterich. “Structure formation and the time dependence of quintessence”. In: *Phys. Rev. D* 64.12 (Dec. 2001), p. 123520. eprint: [arXiv:astro-ph/0107525](#).
- [31] R. Lynds. “The Absorption-Line Spectrum of 4c 05.34”. In: *Astrophys. J. Lett.* 164 (Mar. 1971), pp. L73+.
- [32] P. McDonald. “Toward a Measurement of the Cosmological Geometry at $z \sim 2$: Predicting Ly α Forest Correlation in Three Dimensions and the Potential of Future Data Sets”. In: *Astrophys. J.* 585 (Mar. 2003), pp. 34–51. eprint: [arXiv:astro-ph/0108064](#).
- [33] A. Slosar et al. “The acoustic peak in the Lyman alpha forest”. In: *J. Cosmology Astropart. Phys.* 10 (Oct. 2009), pp. 19–+. arXiv: 0906.2414 [[astro-ph.CO](#)].

- [34] U. Seljak. “Bias, redshift space distortions and primordial nongaussianity of nonlinear transformations: application to Ly- α forest”. In: *J. Cosmology Astropart. Phys.* 3, 004 (Mar. 2012), p. 4. arXiv: 1201.0594 [astro-ph.CO].
- [35] M. White. “The Ly-a forest”. In: *The Davis Meeting On Cosmic Inflation. 2003 March 22-25, Davis CA., p.18*. Mar. 2003.
- [36] P. McDonald and D. J. Eisenstein. “Dark energy and curvature from a future baryonic acoustic oscillation survey using the Lyman- α forest”. In: *Phys. Rev. D* 76.6 (Sept. 2007), pp. 063009–+. eprint: arXiv:astro-ph/0607122.
- [37] M. McQuinn and M. White. “On estimating Ly α forest correlations between multiple sight-lines”. In: *Mon. Not. R. Astron. Soc.* 415 (Aug. 2011), pp. 2257–2269. arXiv: 1102.1752 [astro-ph.CO].
- [38] A. Slosar et al. “The Lyman- α forest in three dimensions: measurements of large scale flux correlations from BOSS 1st-year data”. In: *J. Cosmology Astropart. Phys.* 9 (Sept. 2011), pp. 1–+. arXiv: 1104.5244 [astro-ph.CO].
- [39] N. G. Busca et al. “Baryon acoustic oscillations in the Ly α forest of BOSS quasars”. In: *Astron. Astrophys.* 552, A96 (Apr. 2013), A96. arXiv: 1211.2616 [astro-ph.CO].
- [40] A. Slosar et al. “Measurement of baryon acoustic oscillations in the Lyman- α forest fluctuations in BOSS data release 9”. In: *J. Cosmology Astropart. Phys.* 4, 026 (Apr. 2013), p. 26. arXiv: 1301.3459 [astro-ph.CO].
- [41] D. Kirkby et al. “Fitting methods for baryon acoustic oscillations in the Lyman- α forest fluctuations in BOSS data release 9”. In: *J. Cosmology Astropart. Phys.* 3, 024 (Mar. 2013), p. 24. arXiv: 1301.3456 [astro-ph.CO].
- [42] T. Delubac et al. “Baryon Acoustic Oscillations in the Ly{\alpha} forest of BOSS DR11 quasars”. In: *ArXiv e-prints* (Apr. 2014). arXiv: 1404.1801.
- [43] M. White et al. “Particle Mesh Simulations of the Ly α Forest and the Signature of Baryon Acoustic Oscillations in the Intergalactic Medium”. In: *Astrophys. J.* 713 (Apr. 2010), pp. 383–393. arXiv: 0911.5341.
- [44] M. McQuinn et al. “The signatures of large-scale temperature and intensity fluctuations in the Lyman α forest”. In: *Mon. Not. R. Astron. Soc.* 415 (July 2011), pp. 977–992. arXiv: 1010.5250 [astro-ph.CO].
- [45] M. McQuinn and G. Worseck. “The case against large intensity fluctuations in the $z \sim 2.5$ He II Ly α forest”. In: *Mon. Not. R. Astron. Soc.* 440 (May 2014), pp. 2406–2418. arXiv: 1306.4985.
- [46] A. Pontzen. “Scale-dependent bias in the baryonic-acoustic-oscillation-scale intergalactic neutral hydrogen”. In: *Phys. Rev. D* 89.8, 083010 (Apr. 2014), p. 083010. arXiv: 1402.0506.
- [47] S. Gontcho A Gontcho, J. Miralda-Escudé, and N. G. Busca. “On the effect of the ionizing background on the Ly α forest autocorrelation function”. In: *Mon. Not. R. Astron. Soc.* 442 (July 2014), pp. 187–195.
- [48] M. M. Pieri. “The C IV Forest as a Probe of Baryon Acoustic Oscillations”. In: *ArXiv e-prints* (Apr. 2014). arXiv: 1404.4569.
- [49] P. McDonald and A. Roy. “Clustering of dark matter tracers: generalizing bias for the coming era of precision LSS”. In: *J. Cosmology Astropart. Phys.* 8, 020 (Aug. 2009), p. 20. arXiv: 0902.0991 [astro-ph.CO].

- [50] T. Baldauf et al. “Evidence for quadratic tidal tensor bias from the halo bispectrum”. In: *Phys. Rev. D* 86.8, 083540 (Oct. 2012), p. 083540. arXiv: 1201.4827 [astro-ph.CO].
- [51] P. McDonald. “Clustering of dark matter tracers: Renormalizing the bias parameters”. In: *Phys. Rev. D* 74.10 (Nov. 2006), pp. 103512–+.
- [52] U. Seljak and P. McDonald. “Distribution function approach to redshift space distortions”. In: *J. Cosmology Astropart. Phys.* 11, 039 (Nov. 2011), p. 39. arXiv: 1109.1888 [astro-ph.CO].
- [53] Z. Vlah et al. “Distribution function approach to redshift space distortions. Part IV: perturbation theory applied to dark matter”. In: *J. Cosmology Astropart. Phys.* 11, 009 (Nov. 2012), p. 9. arXiv: 1207.0839 [astro-ph.CO].
- [54] T. Okumura, U. Seljak, and V. Desjacques. “Distribution function approach to redshift space distortions. Part III: halos and galaxies”. In: *J. Cosmology Astropart. Phys.* 11, 014 (Nov. 2012), p. 14. arXiv: 1206.4070 [astro-ph.CO].
- [55] R. Reyes et al. “Confirmation of general relativity on large scales from weak lensing and galaxy velocities”. In: *Nature* 464 (Mar. 2010), pp. 256–258. arXiv: 1003.2185 [astro-ph.CO].
- [56] Y.-S. Song et al. “Complementarity of weak lensing and peculiar velocity measurements in testing general relativity”. In: *Phys. Rev. D* 84.8, 083523 (Oct. 2011), p. 083523. arXiv: 1011.2106 [astro-ph.CO].
- [57] N. Kaiser. “Clustering in real space and in redshift space”. In: *Mon. Not. R. Astron. Soc.* 227 (July 1987), pp. 1–21.
- [58] A. Stril, R. N. Cahn, and E. V. Linder. “Testing standard cosmology with large-scale structure”. In: *Mon. Not. R. Astron. Soc.* 404 (May 2010), pp. 239–246. arXiv: 0910.1833 [astro-ph.CO].
- [59] D. Huterer et al. “Growth of Cosmic Structure: Probing Dark Energy Beyond Expansion”. In: *ArXiv e-prints* (Sept. 2013). arXiv: 1309.5385 [astro-ph.CO].
- [60] N. Hamaus, U. Seljak, and V. Desjacques. “Optimal weighting in galaxy surveys: Application to redshift-space distortions”. In: *Phys. Rev. D* 86.10, 103513 (Nov. 2012), p. 103513. arXiv: 1207.1102 [astro-ph.CO].
- [61] F. C. van den Bosch et al. “Cosmological constraints from a combination of galaxy clustering and lensing - I. Theoretical framework”. In: *Mon. Not. R. Astron. Soc.* 430 (Apr. 2013), pp. 725–746. arXiv: 1206.6890 [astro-ph.CO].
- [62] J. L. Tinker et al. “Cosmological Constraints from Galaxy Clustering and the Mass-to-number Ratio of Galaxy Clusters”. In: *Astrophys. J.* 745, 16 (Jan. 2012), p. 16. arXiv: 1104.1635 [astro-ph.CO].
- [63] J. Yoo and U. Seljak. “Joint analysis of gravitational lensing, clustering, and abundance: Toward the unification of large-scale structure analysis”. In: *Phys. Rev. D* 86.8, 083504 (Oct. 2012), p. 083504. arXiv: 1207.2471 [astro-ph.CO].
- [64] R. Mandelbaum et al. “Cosmological parameter constraints from galaxy-galaxy lensing and galaxy clustering with the SDSS DR7”. In: *Mon. Not. R. Astron. Soc.* 432 (June 2013), pp. 1544–1575. arXiv: 1207.1120 [astro-ph.CO].
- [65] B. Jain et al. “Novel Probes of Gravity and Dark Energy”. In: *ArXiv e-prints* (Sept. 2013). arXiv: 1309.5389 [astro-ph.CO].

- [66] B. Jain, V. Vikram, and J. Sakstein. “Astrophysical Tests of Modified Gravity: Constraints from Distance Indicators in the Nearby Universe”. In: *Astrophys. J.* 779, 39 (Dec. 2013), p. 39. arXiv: 1204.6044 [astro-ph.CO].
- [67] B. A. Reid and M. White. “Towards an accurate model of the redshift-space clustering of haloes in the quasi-linear regime”. In: *Mon. Not. R. Astron. Soc.* 417 (Nov. 2011), pp. 1913–1927. arXiv: 1105.4165 [astro-ph.CO].
- [68] J. Carlson, B. Reid, and M. White. “Convolution Lagrangian perturbation theory for biased tracers”. In: *Mon. Not. R. Astron. Soc.* 429 (Feb. 2013), pp. 1674–1685. arXiv: 1209.0780 [astro-ph.CO].
- [69] Y.-S. Song et al. “Chasing unbiased spectra of the Universe”. In: *Phys. Rev. D* 87.12, 123510 (June 2013), p. 123510. arXiv: 1301.3133 [astro-ph.CO].
- [70] B. A. Reid and M. White. “Towards an accurate model of the redshift-space clustering of haloes in the quasi-linear regime”. In: *Mon. Not. R. Astron. Soc.* 417 (Nov. 2011), pp. 1913–1927. arXiv: 1105.4165 [astro-ph.CO].
- [71] Z. Vlah et al. “Distribution function approach to redshift space distortions. Part V: perturbation theory applied to dark matter halos”. In: *ArXiv e-prints* (Aug. 2013). arXiv: 1308.6294 [astro-ph.CO].
- [72] L. Samushia, W. J. Percival, and A. Raccanelli. “Interpreting large-scale redshift-space distortion measurements”. In: *Mon. Not. R. Astron. Soc.* 420 (Mar. 2012), pp. 2102–2119. arXiv: 1102.1014 [astro-ph.CO].
- [73] T. Okumura et al. “Large-Scale Anisotropic Correlation Function of SDSS Luminous Red Galaxies”. In: *Astrophys. J.* 676 (Apr. 2008), pp. 889–898. arXiv: 0711.3640.
- [74] Y.-S. Song et al. “Measuring coherent motions in the universe”. In: *J. Cosmology Astropart. Phys.* 5, 020 (May 2011), p. 20. arXiv: 1006.4630 [astro-ph.CO].
- [75] F. Beutler et al. “The 6dF Galaxy Survey: $z = 0$ measurements of the growth rate and σ_8 ”. In: *Mon. Not. R. Astron. Soc.* 423 (July 2012), pp. 3430–3444. arXiv: 1204.4725 [astro-ph.CO].
- [76] Y.-S. Song et al. “Cosmological Tests using Redshift Space Clustering in BOSS DR11”. In: *ArXiv e-prints* (July 2014). arXiv: 1407.2257.
- [77] C. Blake et al. “The WiggleZ Dark Energy Survey: the growth rate of cosmic structure since redshift $z=0.9$ ”. In: *Mon. Not. R. Astron. Soc.* 415 (Aug. 2011), pp. 2876–2891. arXiv: 1104.2948 [astro-ph.CO].
- [78] B. A. Reid et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: measurements of the growth of structure and expansion rate at $z = 0.57$ from anisotropic clustering”. In: *Mon. Not. R. Astron. Soc.* 426 (Nov. 2012), pp. 2719–2737. arXiv: 1203.6641 [astro-ph.CO].
- [79] F. Beutler et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Testing gravity with redshift-space distortions using the power spectrum multipoles”. In: *ArXiv:132.4611 (submitted)* (Dec. 2013). arXiv: 1312.4611.
- [80] L. Samushia et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: measuring growth rate and geometry with anisotropic clustering”. In: *Mon. Not. R. Astron. Soc.* 439 (Apr. 2014), pp. 3504–3519. arXiv: 1312.4899.

- [81] E. A. Kazin et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: measuring $H(z)$ and $D_A(z)$ at $z = 0.57$ with clustering wedges”. In: *Mon.-Not. R. Astron. Soc.* (Aug. 2013). arXiv: 1303.4391 [astro-ph.CO].
- [82] A. G. Sánchez et al. “The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological implications of the full shape of the clustering wedges in the data release 10 and 11 galaxy samples”. In: *Mon. Not. R. Astron. Soc.* 440 (May 2014), pp. 2692–2713. arXiv: 1312.4854 [astro-ph.CO].
- [83] W. J. Percival et al. “The clustering of Galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: including covariance matrix errors”. In: *Mon. Not. R. Astron. Soc.* 439 (Apr. 2014), pp. 2531–2541. arXiv: 1312.4841 [astro-ph.CO].
- [84] A. Font-Ribera et al. “DESI and other Dark Energy experiments in the era of neutrino mass measurements”. In: *J. Cosmology Astropart. Phys.* 5, 023 (May 2014), p. 23. arXiv: 1308.4164.
- [85] L. Samushia et al. “The clustering of galaxies in the SDSS-III DR9 Baryon Oscillation Spectroscopic Survey: testing deviations from Λ and general relativity using anisotropic clustering of galaxies”. In: *Mon. Not. R. Astron. Soc.* 429 (Feb. 2013), pp. 1514–1528. arXiv: 1206.5309 [astro-ph.CO].
- [86] W. J. Percival et al. “The 2dF Galaxy Redshift Survey: spherical harmonics analysis of fluctuations in the final catalogue”. In: *Mon. Not. R. Astron. Soc.* 353 (Oct. 2004), pp. 1201–1218. eprint: arXiv:astro-ph/0406513.
- [87] L. Guzzo et al. “A test of the nature of cosmic acceleration using galaxy redshift distortions”. In: *Nature* 451 (Jan. 2008), pp. 541–544. arXiv: 0802.1944.
- [88] M. J. Drinkwater et al. “The WiggleZ Dark Energy Survey: survey design and first data release”. In: *Mon. Not. R. Astron. Soc.* 401 (Jan. 2010), pp. 1429–1452. arXiv: 0911.4246 [astro-ph.CO].
- [89] E. Komatsu et al. “Seven-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation”. In: *Astrophys. J. Supp.* 192, 18 (Feb. 2011), p. 18. arXiv: 1001.4538 [astro-ph.CO].
- [90] C. Alcock and B. Paczynski. “An evolution free test for non-zero cosmological constant”. In: *Nature* 281 (Oct. 1979), pp. 358–+.
- [91] P. McDonald. “Toward a Measurement of the Cosmological Geometry at $z \sim 2$: Predicting Ly α Forest Correlation in Three Dimensions and the Potential of Future Data Sets”. In: *Astrophys. J.* 585 (Mar. 2003), pp. 34–51. eprint: arXiv:astro-ph/0108064.
- [92] H.-J. Seo and D. J. Eisenstein. “Improved Forecasts for the Baryon Acoustic Oscillations and Cosmological Distance Scale”. In: *Astrophys. J.* 665 (Aug. 2007), pp. 14–24. eprint: arXiv:astro-ph/0701079.
- [93] H. A. Feldman, N. Kaiser, and J. A. Peacock. “Power-spectrum analysis of three-dimensional redshift surveys”. In: *Astrophys. J.* 426 (May 1994), pp. 23–37.
- [94] N. Mostek et al. “The DEEP2 Galaxy Redshift Survey: Clustering Dependence on Galaxy Stellar Mass and Star Formation Rate at $z \sim 1$ ”. In: *Astrophys. J.* 767, 89 (Apr. 2013), p. 89. arXiv: 1210.6694 [astro-ph.CO].
- [95] N. P. Ross et al. “Clustering of Low-redshift ($z \leq 2.2$) Quasars from the Sloan Digital Sky Survey”. In: *Astrophys. J.* 697 (June 2009), pp. 1634–1655. arXiv: 0903.3230 [astro-ph.CO].

- [96] N. Mostek et al. “Mapping the universe with BigBOSS”. In: *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*. Vol. 8446. Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series. Sept. 2012.
- [97] R. Laureijs et al. “Euclid Definition Study Report”. In: *ArXiv e-prints* (Oct. 2011). arXiv: 1110.3193 [astro-ph.CO].
- [98] D. Spergel et al. “Wide-Field InfraRed Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA Final Report”. In: *ArXiv e-prints* (May 2013). arXiv: 1305.5422 [astro-ph.IM].
- [99] A. G. Riess et al. “A 3% Solution: Determination of the Hubble Constant with the Hubble Space Telescope and Wide Field Camera 3”. In: *Astrophys. J.* 730, 119 (Apr. 2011), p. 119. arXiv: 1103.2976 [astro-ph.CO].
- [100] X. Xu et al. “Measuring D_A and H at $z=0.35$ from the SDSS DR7 LRGs using baryon acoustic oscillations”. In: *Mon. Not. R. Astron. Soc.* 431 (May 2013), pp. 2834–2860. arXiv: 1206.6732 [astro-ph.CO].
- [101] A. Font-Ribera et al. “Quasar-Lyman α forest cross-correlation from BOSS DR11: Baryon Acoustic Oscillations”. In: *J. Cosmology Astropart. Phys.* 5, 027 (May 2014), p. 27. arXiv: 1311.1767.
- [102] M. Sullivan et al. “SNLS3: Constraints on Dark Energy Combining the Supernova Legacy Survey Three-year Data with Other Probes”. In: *Astrophys. J.* 737, 102 (Aug. 2011), p. 102. arXiv: 1104.1444 [astro-ph.CO].
- [103] M. White. “Shot noise and reconstruction of the acoustic peak”. In: *ArXiv e-prints* (Apr. 2010). arXiv: 1004.0250 [astro-ph.CO].
- [104] P. McDonald et al. “The Ly α Forest Power Spectrum from the Sloan Digital Sky Survey”. In: *Astrophys. J. Supp.* 163 (Mar. 2006), pp. 80–109. eprint: arXiv:astro-ph/0405013.
- [105] P. McDonald et al. “The Linear Theory Power Spectrum from the Ly α Forest in the Sloan Digital Sky Survey”. In: *Astrophys. J.* 635 (Dec. 2005), pp. 761–783.
- [106] P. McDonald et al. “Physical effects on the Ly α forest flux power spectrum: damping wings, ionizing radiation fluctuations and galactic winds”. In: *Mon. Not. R. Astron. Soc.* 360 (July 2005), pp. 1471–1482.
- [107] R. Mandelbaum et al. “Precision cosmology from the Lyman- α forest: power spectrum and bispectrum”. In: *Mon. Not. R. Astron. Soc.* 344 (Sept. 2003), pp. 776–788.
- [108] P. McDonald et al. “A Measurement of the Temperature-Density Relation in the Intergalactic Medium Using a New Ly α Absorption-Line Fitting Method”. In: *Astrophys. J.* 562 (Nov. 2001), pp. 52–75.
- [109] P. McDonald et al. “The Observed Probability Distribution Function, Power Spectrum, and Correlation Function of the Transmitted Flux in the Ly α Forest”. In: *Astrophys. J.* 543 (Nov. 2000), pp. 1–23.
- [110] T.-S. Kim et al. “The power spectrum of the flux distribution in the Lyman α forest of a large sample of UVES QSO absorption spectra (LUQAS)”. In: *Mon. Not. R. Astron. Soc.* 347 (Jan. 2004), pp. 355–366. eprint: arXiv:astro-ph/0308103.
- [111] V. Iršič et al. “Detection of Ly- β auto-correlations and Ly- α -Ly- β cross-correlations in BOSS Data Release 9”. In: *ArXiv e-prints* (July 2013). arXiv: 1307.3403 [astro-ph.CO].

- [112] A. Font-Ribera et al. “The large-scale quasar-Lyman α forest cross-correlation from BOSS”. In: *J. Cosmology Astropart. Phys.* 5, 018 (May 2013), p. 18. arXiv: 1303.1937 [astro-ph.CO].
- [113] N. Palanque-Delabrouille et al. “The one-dimensional Ly-alpha forest power spectrum from BOSS”. In: *ArXiv e-prints* (June 2013). arXiv: 1306.5896 [astro-ph.CO].
- [114] U. Seljak, A. Slosar, and P. McDonald. “Cosmological parameters from combining the Lyman- α forest with CMB, galaxy clustering and SN constraints”. In: *Journal of Cosmology and Astro-Particle Physics* 10 (Oct. 2006), pp. 14–+.
- [115] U. Seljak et al. “Cosmological parameter analysis including SDSS Ly α forest and galaxy bias: Constraints on the primordial spectrum of fluctuations, neutrino mass, and dark energy”. In: *Phys. Rev. D* 71.10 (May 2005), pp. 103515–+.
- [116] A. Kosowsky and M. S. Turner. “CBR anisotropy and the running of the scalar spectral index”. In: *Phys. Rev. D* 52 (Aug. 1995), p. 1739. eprint: arXiv:astro-ph/9504071.
- [117] S. Dodelson. *Modern cosmology.*, 2003.
- [118] P.A.R. Ade et al. “Planck 2015 results. XX. Constraints on inflation”. In: *ArXiv e-prints* (2015). arXiv: 1502.02114 [astro-ph.CO].
- [119] A. R. Liddle and D. H. Lyth. *Cosmological Inflation and Large-Scale Structure.*, June 2000.
- [120] R. Easther and H. V. Peiris. “Implications of a running spectral index for slow roll inflation”. In: *J. Cosmology Astropart. Phys.* 9, 010 (Sept. 2006), p. 10. eprint: arXiv:astro-ph/0604214.
- [121] D. J. Chung, G. Shiu, and M. Trodden. “Running of the scalar spectral index from inflationary models”. In: *Phys. Rev. D* 68.6, 063501 (Sept. 2003), p. 063501. eprint: arXiv:astro-ph/0305193.
- [122] P.A.R. Ade et al. “Planck 2015 results. XVII. Constraints on primordial non-Gaussianity”. In: *ArXiv e-prints* (2015). arXiv: 1502.01592 [astro-ph.CO].
- [123] Xingang Chen et al. “Observational signatures and non-Gaussianities of general single field inflation”. In: *JCAP* 0701 (2007), p. 002. arXiv: hep-th/0605045 [hep-th].
- [124] Justin Khoury and Federico Piazza. “Rapidly-Varying Speed of Sound, Scale Invariance and Non-Gaussian Signatures”. In: *JCAP* 0907 (2009), p. 026. arXiv: 0811.3633 [hep-th].
- [125] Johannes Noller and Joao Magueijo. “Non-Gaussianity in single field models without slow-roll”. In: *Phys.Rev.* D83 (2011), p. 103511. arXiv: 1102.0275 [astro-ph.CO].
- [126] Raquel H. Ribeiro. “Inflationary signatures of single-field models beyond slow-roll”. In: *JCAP* 1205 (2012), p. 037. arXiv: 1202.4453 [astro-ph.CO].
- [127] Johannes Noller. “Constraining fast-roll inflation”. In: *ArXiv e-prints* (2012). arXiv: 1205.5796 [astro-ph.CO].
- [128] Xingang Chen and Yi Wang. “Large non-Gaussianities with Intermediate Shapes from Quasi-Single Field Inflation”. In: *Phys.Rev.* D81 (2010), p. 063511. arXiv: 0909.0496 [astro-ph.CO].
- [129] Xingang Chen and Yi Wang. “Quasi-Single Field Inflation and Non-Gaussianities”. In: *JCAP* 1004 (2010), p. 027. arXiv: 0911.3380 [hep-th].
- [130] Sujata Gupta et al. “Non-Gaussian signatures in the cosmic background radiation from warm inflation”. In: *Phys.Rev.* D66 (2002), p. 043510. arXiv: astro-ph/0205152 [astro-ph].

- [131] Ian G Moss and Chun Xiong. “Non-Gaussianity in fluctuations from warm inflation”. In: *JCAP* 0704 (2007), p. 007. arXiv: [astro-ph/0701302](#) [[astro-ph](#)].
- [132] R. Holman and Andrew J. Tolley. “Enhanced Non-Gaussianity from Excited Initial States”. In: *JCAP* 0805 (2008), p. 001. arXiv: [0710.1302](#) [[hep-th](#)].
- [133] Pieter Daniel Meerburg, Jan Pieter van der Schaar, and Pier Stefano Corasaniti. “Signatures of Initial State Modifications on Bispectrum Statistics”. In: *JCAP* 0905 (2009), p. 018. arXiv: [0901.4044](#) [[hep-th](#)].
- [134] Nishant Agarwal et al. “Effective field theory and non-Gaussianity from general inflationary states”. In: *ArXiv e-prints* (2012). arXiv: [1212.1172](#) [[hep-th](#)].
- [135] David Langlois et al. “Primordial perturbations and non-Gaussianities in DBI and general multi-field inflation”. In: *Phys.Rev. D* 78 (2008), p. 063523. arXiv: [0806.0336](#) [[hep-th](#)].
- [136] Frederico Arroja, Shuntaro Mizuno, and Kazuya Koyama. “Non-gaussianity from the bispectrum in general multiple field inflation”. In: *JCAP* 0808 (2008), p. 015. arXiv: [0806.0619](#) [[astro-ph](#)].
- [137] Sebastien Renaux-Petel. “Combined local and equilateral non-Gaussianities from multifield DBI inflation”. In: *JCAP* 0910 (2009), p. 012. arXiv: [0907.2476](#) [[hep-th](#)].
- [138] David Langlois and Angela Lepidi. “General treatment of isocurvature perturbations and non-Gaussianities”. In: *JCAP* 1101 (2011), p. 008. arXiv: [1007.5498](#) [[astro-ph.CO](#)].
- [139] David Langlois and Bartjan van Tent. “Hunting for Isocurvature Modes in the CMB non-Gaussianities”. In: *Class.Quant.Grav.* 28 (2011), p. 222001. arXiv: [1104.2567](#) [[astro-ph.CO](#)].
- [140] David Langlois and Bartjan van Tent. “Isocurvature modes in the CMB bispectrum”. In: *JCAP* 1207 (2012), p. 040. arXiv: [1204.5042](#) [[astro-ph.CO](#)].
- [141] E. Komatsu. “The Pursuit of Non-Gaussian Fluctuations in the Cosmic Microwave Background”. In: *ArXiv Astrophysics e-prints* (June 2002). eprint: [arXiv:astro-ph/0206039](#).
- [142] C. L. Bennett et al. “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results”. In: *ArXiv e-prints* (Dec. 2012). arXiv: [1212.5225](#) [[astro-ph.CO](#)].
- [143] Planck Collaboration et al. “Planck 2013 Results. XXIV. Constraints on primordial non-Gaussianity”. In: *ArXiv e-prints* (Mar. 2013). arXiv: [1303.5084](#) [[astro-ph.CO](#)].
- [144] N. Dalal et al. “Imprints of primordial non-Gaussianities on large-scale structure: Scale-dependent bias and abundance of virialized objects”. In: *Phys. Rev. D* 77.12 (June 2008), pp. 123514–+. arXiv: [0710.4560](#).
- [145] A. Slosar et al. “Constraints on local primordial non-Gaussianity from large scale structure”. In: *Journal of Cosmology and Astro-Particle Physics* 8 (Aug. 2008), pp. 31–+. arXiv: [0805.3580](#).
- [146] S. Ho et al. “Sloan Digital Sky Survey III Photometric Quasar Clustering: Probing the Initial Conditions of the Universe using the Largest Volume”. In: *ArXiv e-prints* (Nov. 2013). arXiv: [1311.2597](#) [[astro-ph.CO](#)].
- [147] T. Giannantonio et al. “Improved primordial non-Gaussianity constraints from measurements of galaxy clustering and the integrated Sachs-Wolfe effect”. In: *Phys. Rev. D* 89.2, 023511 (Jan. 2014), p. 023511. arXiv: [1303.1349](#) [[astro-ph.CO](#)].
- [148] N. Agarwal et al. “Characterizing unknown systematics in large scale structure surveys”. In: *ArXiv e-prints* (Sept. 2013). arXiv: [1309.2954](#) [[astro-ph.CO](#)].

- [149] C. Pitrou, J.-P. Uzan, and F. Bernardeau. “The cosmic microwave background bispectrum from the non-linear evolution of the cosmological perturbations”. In: *J. Cosmology-Astropart. Phys.* 7, 003 (July 2010), p. 3. arXiv: 1003.0481 [astro-ph.CO].
- [150] N. Bartolo, S. Matarrese, and A. Riotto. “Non-Gaussianity and the Cosmic Microwave Background Anisotropies”. In: *Advances in Astronomy 2010* (2010). arXiv: 1001.3957 [astro-ph.CO].
- [151] A. Becker and D. Huterer. “First Constraints on the Running of Non-Gaussianity”. In: *Physical Review Letters* 109.12, 121302 (Sept. 2012), p. 121302. arXiv: 1207.5788 [astro-ph.CO].
- [152] A. Becker, D. Huterer, and K. Kadota. “Constraining scale-dependent non-Gaussianity with future large-scale structure and the CMB”. In: *J. Cosmology Astropart. Phys.* 12, 034 (Dec. 2012), p. 34. arXiv: 1206.6165 [astro-ph.CO].
- [153] U. Seljak. “Extracting Primordial Non-Gaussianity without Cosmic Variance”. In: *Physical Review Letters* 102.2, 021302 (Jan. 2009), p. 021302. arXiv: 0807.1770.
- [154] P. McDonald and U. Seljak. “How to evade the sample variance limit on measurements of redshift-space distortions”. In: *J. Cosmology Astropart. Phys.* 10, 007 (Oct. 2009), p. 7. arXiv: 0810.0323.
- [155] V. Desjacques, U. Seljak, and I. T. Iliev. “Scale-dependent bias induced by local non-Gaussianity: a comparison to N-body simulations”. In: *Mon. Not. R. Astron. Soc.* 396 (June 2009), pp. 85–96. arXiv: 0811.2748.
- [156] D. Huterer, C. E. Cunha, and W. Fang. “Calibration errors unleashed: effects on cosmological parameters and requirements for large-scale structure surveys”. In: *Mon. Not. R. Astron. Soc.* 432 (July 2013), pp. 2945–2961. arXiv: 1211.1015 [astro-ph.CO].
- [157] B. Leistedt et al. “Estimating the large-scale angular power spectrum in the presence of systematics: a case study of Sloan Digital Sky Survey quasars”. In: *Mon. Not. R. Astron. Soc.* (Sept. 2013). arXiv: 1306.0005 [astro-ph.CO].
- [158] J. Lesgourgues et al. *Neutrino Cosmology.*, Feb. 2013.
- [159] Gianpiero Mangano et al. “Relic neutrino decoupling including flavor oscillations”. In: *Nucl.-Phys.* B729 (2005), pp. 221–234. arXiv: hep-ph/0506164 [hep-ph].
- [160] J. Hamann et al. “Cosmology seeking friendship with sterile neutrinos”. In: *Nuclear Physics B Proceedings Supplements* 217 (Aug. 2011), pp. 72–74.
- [161] W. Fischler and J. Meyers. “Dark radiation emerging after big bang nucleosynthesis?” In: *Phys. Rev. D* 83.6, 063520 (Mar. 2011), p. 063520. arXiv: 1011.3501 [astro-ph.CO].
- [162] S. Bashinsky and U. Seljak. “Signatures of relativistic neutrinos in CMB anisotropy and matter clustering”. In: *Phys. Rev. D* 69.8, 083002 (Apr. 2004), p. 083002. eprint: arXiv: astro-ph/0310198.
- [163] E. Calabrese et al. “Limits on dark radiation, early dark energy, and relativistic degrees of freedom”. In: *Phys. Rev. D* 83.12, 123504 (June 2011), p. 123504. arXiv: 1103.4132 [astro-ph.CO].
- [164] S. Ho et al. “Correlation of CMB with large-scale structure. I. Integrated Sachs-Wolfe tomography and cosmological implications”. In: *Phys. Rev. D* 78.4, 043519 (Aug. 2008), p. 043519. arXiv: 0801.0642.

- [165] P. Zhang et al. “Probing Gravity at Cosmological Scales by Measurements which Test the Relationship between Gravitational Lensing and Matter Overdensity”. In: *Physical Review Letters* 99.14 (Oct. 2007), pp. 141302–+. arXiv: 0704.1932.
- [166] J.R. Fergusson, M. Liguori, and E.P.S. Shellard. “General CMB and Primordial Bispectrum Estimation I: Mode Expansion, Map-Making and Measures of f_{NL} ”. In: *Phys.Rev.* D82 (2010), p. 023502. arXiv: 0912.5516 [astro-ph.CO].
- [167] M. Liguori et al. “Primordial non-Gaussianity and Bispectrum Measurements in the Cosmic Microwave Background and Large-Scale Structure”. In: *Adv.Astron.* 2010 (2010), p. 980523. arXiv: 1001.4707 [astro-ph.CO].
- [168] P. Zhang. “The dark flow induced small-scale kinetic Sunyaev-Zel’dovich effect”. In: *Mon.-Not. R. Astron. Soc.* 407 (Sept. 2010), pp. L36–L40. arXiv: 1004.0990 [astro-ph.CO].
- [169] P. Zhang and A. Stebbins. “Confirmation of the Copernican Principle at Gpc Radial Scale and above from the Kinetic Sunyaev-Zel’dovich Effect Power Spectrum”. In: *Physical Review Letters* 107.4, 041301 (July 2011), p. 041301. arXiv: 1009.3967 [astro-ph.CO].
- [170] Z. Li, P. Zhang, and X. Chen. “Anomalous Anisotropic Cross-correlations between WMAP CMB Maps and SDSS Galaxy Distribution and Implications on the Dark Flow Scenario”. In: *Astrophys. J.* 758, 130 (Oct. 2012), p. 130. arXiv: 1209.0520 [astro-ph.CO].
- [171] S. Ho, S. Dedeo, and D. Spergel. “Finding the Missing Baryons Using CMB as a Backlight”. In: *ArXiv e-prints* (Mar. 2009). arXiv: 0903.2845 [astro-ph.CO].
- [172] J. A. Newman. “Calibrating Redshift Distributions beyond Spectroscopic Limits with Cross-Correlations”. In: *Astrophys. J.* 684 (Sept. 2008), pp. 88–101. arXiv: 0805.1409.
- [173] M. McQuinn and M. White. “On using angular cross-correlations to determine source redshift distributions”. In: *Mon. Not. R. Astron. Soc.* 433 (Aug. 2013), pp. 2857–2883. arXiv: 1302.0857 [astro-ph.CO].
- [174] S. J. Schmidt et al. “Recovering redshift distributions with cross-correlations: pushing the boundaries”. In: *Mon. Not. R. Astron. Soc.* 431 (June 2013), pp. 3307–3318. arXiv: 1303.0292 [astro-ph.CO].
- [175] J. Coupon, T. Broadhurst, and K. Umetsu. “Cluster Lensing Profiles Derived from a Redshift Enhancement of Magnified BOSS-survey Galaxies”. In: *Astrophys. J.* 772, 65 (July 2013), p. 65. arXiv: 1303.6588 [astro-ph.CO].
- [176] M. Cacciato et al. “Cosmological constraints from a combination of galaxy clustering and lensing - III. Application to SDSS data”. In: *Mon. Not. R. Astron. Soc.* 430 (Apr. 2013), pp. 767–786. arXiv: 1207.0503 [astro-ph.CO].
- [177] Y.-C. Cai and G. Bernstein. “Combining weak-lensing tomography and spectroscopic redshift surveys”. In: *Mon. Not. R. Astron. Soc.* 422 (May 2012), pp. 1045–1056. arXiv: 1112.4478 [astro-ph.CO].
- [178] E. Gaztañaga et al. “Cross-correlation of spectroscopic and photometric galaxy surveys: cosmology from lensing and redshift distortions”. In: *Mon. Not. R. Astron. Soc.* 422 (June 2012), pp. 2904–2930. arXiv: 1109.4852 [astro-ph.CO].
- [179] D. Kirk et al. “Optimising Spectroscopic and Photometric Galaxy Surveys: Same-sky Benefits for Dark Energy and Modified Gravity”. In: *ArXiv e-prints* (July 2013). arXiv: 1307.8062 [astro-ph.CO].

- [180] R. de Putter, O. Doré, and M. Takada. “The Synergy between Weak Lensing and Galaxy Redshift Surveys”. In: *ArXiv e-prints* (Aug. 2013). arXiv: 1308.6070 [astro-ph.CO].
- [181] T. Y. Lam et al. “Testing Gravity with the Stacked Phase Space around Galaxy Clusters”. In: *Physical Review Letters* 109.5, 051301 (Aug. 2012), p. 051301. arXiv: 1202.4501 [astro-ph.CO].
- [182] R. de Putter, O. Doré, and M. Takada. “The Synergy between Weak Lensing and Galaxy Redshift Surveys”. In: *ArXiv e-prints* (Aug. 2013). arXiv: 1308.6070 [astro-ph.CO].
- [183] M. Betoule et al. “Improved cosmological constraints from a joint analysis of the SDSS-II and SNLS supernova samples”. In: *Astron. Astrophys.* 568, A22 (Aug. 2014), A22. arXiv: 1401.4064.
- [184] C. Conroy, R. H. Wechsler, and A. V. Kravtsov. “Modeling Luminosity-dependent Galaxy Clustering through Cosmic Time”. In: *Astrophys. J.* 647 (Aug. 2006), pp. 201–214. eprint: arXiv:astro-ph/0512234.
- [185] D. J. Eisenstein et al. “Spectroscopic Target Selection for the Sloan Digital Sky Survey: The Luminous Red Galaxy Sample”. In: *Astron. J.* 122 (Nov. 2001), pp. 2267–2280. eprint: arXiv:astro-ph/0108153.
- [186] D. J. Eisenstein et al. “Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies”. In: *Astrophys. J.* 633 (Nov. 2005), pp. 560–574. eprint: arXiv:astro-ph/0501171.
- [187] S. Ho et al. “Luminous Red Galaxy Population in Clusters at $0.2 \leq z \leq 0.6$ ”. In: *Astrophys. J.* 697 (June 2009), pp. 1358–1368. arXiv: 0706.0727.
- [188] E. A. Kazin et al. “The Baryonic Acoustic Feature and Large-Scale Clustering in the Sloan Digital Sky Survey Luminous Red Galaxy Sample”. In: *Astrophys. J.* 710 (Feb. 2010), pp. 1444–1461. arXiv: 0908.2598 [astro-ph.CO].
- [189] N. Padmanabhan et al. “The clustering of luminous red galaxies in the Sloan Digital Sky Survey imaging data”. In: *Mon. Not. R. Astron. Soc.* 378 (July 2007), pp. 852–872. eprint: arXiv:astro-ph/0605302.
- [190] T. L. John. “Continuous absorption by the negative hydrogen ion reconsidered”. In: *Astron.-Astrophys.* 193 (Mar. 1988), pp. 189–192.
- [191] M. Sawicki. “The 1.6 Micron Bump as a Photometric Redshift Indicator”. In: *Astron. J.* 124 (Dec. 2002), pp. 3050–3060. eprint: arXiv:astro-ph/0209437.
- [192] M. J. I. Brown et al. “An Atlas of Galaxy Spectral Energy Distributions from the Ultraviolet to the Mid-infrared”. In: *Astrophys. J. Supp.* 212, 18 (June 2014), p. 18. arXiv: 1312.3029 [astro-ph.CO].
- [193] S. D. J. Gwyn. “MegaPipe: The MegaCam Image Stacking Pipeline at the Canadian Astronomical Data Centre”. In: *Proc. Astron. Soc. Pacific* 120 (Feb. 2008), pp. 212–223. arXiv: 0710.0370.
- [194] R. M. Cutri et al. *Explanatory Supplement to the WISE All-Sky Data Release Products*. Tech. rep. WISE, Mar. 2012, p. 1.
- [195] J. A. Newman et al. “The DEEP2 Galaxy Redshift Survey: Design, Observations, Data Reduction, and Redshifts”. In: *Astrophys. J. Supp.* 208, 5 (Sept. 2013), p. 5. arXiv: 1203.3192 [astro-ph.CO].

- [196] O. Ilbert et al. “Cosmos Photometric Redshifts with 30-Bands for 2-deg²”. In: *Astrophys. J.* 690 (Jan. 2009), pp. 1236–1249. arXiv: 0809.2101.
- [197] M. White et al. “The Clustering of Massive Galaxies at $z \sim 0.5$ from the First Semester of BOSS Data”. In: *Astrophys. J.* 728, 126 (Feb. 2011), p. 126. arXiv: 1010.4915 [astro-ph.CO].
- [198] A. M. Hopkins and J. F. Beacom. “On the Normalization of the Cosmic Star Formation History”. In: *Astrophys. J.* 651 (Nov. 2006), pp. 142–154. eprint: arXiv:astro-ph/0601463.
- [199] G. Zhu, J. Moustakas, and M. R. Blanton. “The [O II] $\lambda 3727$ Luminosity Function at $z \sim 1$ ”. In: *Astrophys. J.* 701 (Aug. 2009), pp. 86–93. arXiv: 0811.3035.
- [200] W. Rujopakarn et al. “The Evolution of the Star Formation Rate of Galaxies at $0.0 \leq z \leq 1.2$ ”. In: *Astrophys. J.* 718 (Aug. 2010), pp. 1171–1185. eprint: 1006.4359 (astro-ph.CO).
- [201] J. Comparat et al. “Measuring galaxy [OII] emission line doublet with future ground-based wide-field spectroscopic surveys”. In: *ArXiv e-prints* (Oct. 2013). arXiv: 1310.0615 [astro-ph.IM].
- [202] D. J. Matthews et al. “Extended Photometry for the DEEP2 Galaxy Redshift Survey: A Testbed for Photometric Redshift Experiments”. In: *Astrophys. J. Supp.* 204, 21 (Feb. 2013), p. 21. arXiv: 1210.2405 [astro-ph.CO].
- [203] M. Davis et al. “Science Objectives and Early Results of the DEEP2 Redshift Survey”. In: *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*. Ed. by P. Guhathakurta. Vol. 4834. Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series. Feb. 2003, pp. 161–172. eprint: arXiv:astro-ph/0209419.
- [204] J. Comparat et al. “Investigating emission-line galaxy surveys with the Sloan Digital Sky Survey infrastructure”. In: *Mon. Not. R. Astron. Soc.* 428 (Jan. 2013), pp. 1498–1517. arXiv: 1207.4321 [astro-ph.CO].
- [205] R. C. Kennicutt Jr. “Star Formation in Galaxies Along the Hubble Sequence”. In: *Annu.-Rev. Astron. Astrophys.* 36 (1998), pp. 189–232. eprint: arXiv:astro-ph/9807187.
- [206] R. C. Kennicutt and N. J. Evans. “Star Formation in the Milky Way and Nearby Galaxies”. In: *Annu. Rev. Astron. Astrophys.* 50 (Sept. 2012), pp. 531–608. arXiv: 1204.3552 [astro-ph.GA].
- [207] O. Ilbert et al. “Mass assembly in quiescent and star-forming galaxies since $z \simeq 4$ from Ultra-VISTA”. In: *Astron. Astrophys.* 556, A55 (Aug. 2013), A55. arXiv: 1301.3157 [astro-ph.CO].
- [208] O. Le Fèvre et al. “The VIMOS VLT Deep Survey final data release: a spectroscopic sample of 35 016 galaxies and AGN out to $z \sim 6.7$ selected with $17.5 \leq i_{AB} \leq 24.75$ ”. In: *Astron.-Astrophys.* 559, A14 (Nov. 2013), A14. arXiv: 1307.0545 [astro-ph.CO].
- [209] A. L. Coil et al. “The DEEP2 Galaxy Redshift Survey: Color and Luminosity Dependence of Galaxy Clustering at $z \sim 1$ ”. In: *Astrophys. J.* 672 (Jan. 2008), pp. 153–176. arXiv: 0708.0004.
- [210] N. Mostek et al. “The DEEP2 Galaxy Redshift Survey: Clustering Dependence on Galaxy Stellar Mass and Star Formation Rate at $z \sim 1$ ”. In: *Astrophys. J.* 767, 89 (Apr. 2013), p. 89. arXiv: 1210.6694 [astro-ph.CO].
- [211] R. E. Smith et al. “Stable clustering, the halo model and non-linear cosmological power spectra”. In: *Mon. Not. R. Astron. Soc.* 341 (June 2003), pp. 1311–1332. eprint: arXiv: astro-ph/0207664.

- [212] J. E. Geach et al. “HiZELS: a high-redshift survey of H α emitters - I. The cosmic star formation rate and clustering at $z = 2.23$ ”. In: *Mon. Not. R. Astron. Soc.* 388 (Aug. 2008), pp. 1473–1486. arXiv: 0805.2861.
- [213] C. Blake et al. “The WiggleZ Dark Energy Survey: small-scale clustering of Lyman-break galaxies at $z \leq 1$ ”. In: *Mon. Not. R. Astron. Soc.* 395 (May 2009), pp. 240–254. arXiv: 0901.2587 [astro-ph.CO].
- [214] M. Sumiyoshi et al. “Photometric H alpha and [O II] Luminosity Function of SDF and SXDF Galaxies: Implications for Future Baryon Oscillation Surveys”. In: *ArXiv e-prints* (Feb. 2009). arXiv: 0902.2064.
- [215] G. T. Richards et al. “Efficient Photometric Selection of Quasars from the Sloan Digital Sky Survey. II. $\sim 1,000,000$ Quasars from Data Release 6”. In: *Astrophys. J. Supp.* 180 (Jan. 2009), pp. 67–83. arXiv: 0809.3952.
- [216] N. Palanque-Delabrouille et al. “Luminosity function from dedicated SDSS-III and MMT data of quasars in $0.7 < z < 4.0$ selected with a new approach”. In: *Astron. Astrophys.* 551, A29 (Mar. 2013), A29. arXiv: 1209.3968 [astro-ph.CO].
- [217] N. Palanque-Delabrouille et al. “Variability selected high-redshift quasars on SDSS Stripe 82”. In: *Astron. Astrophys.* 530, A122 (June 2011), A122. arXiv: 1012.2391 [astro-ph.CO].
- [218] L. Jiang et al. “A Spectroscopic Survey of Faint Quasars in the SDSS Deep Stripe. I. Preliminary Results from the Co-added Catalog”. In: *Astron. J.* 131 (June 2006), pp. 2788–2800. eprint: arXiv:astro-ph/0602569.
- [219] P. F. Hopkins, G. T. Richards, and L. Hernquist. “An Observational Determination of the Bolometric Quasar Luminosity Function”. In: *Astrophys. J.* 654 (Jan. 2007), pp. 731–753. eprint: arXiv:astro-ph/0605678.
- [220] LSST Science Collaboration et al. “LSST Science Book, Version 2.0”. In: *ArXiv e-prints* (Dec. 2009). arXiv: 0912.0201 [astro-ph.IM].
- [221] D. Stern et al. “Mid-Infrared Selection of Active Galaxies”. In: *Astrophys. J.* 631 (Sept. 2005), pp. 163–168. eprint: arXiv:astro-ph/0410523.
- [222] C. Yèche et al. “Artificial neural networks for quasar selection and photometric redshift determination”. In: *Astron. Astrophys.* 523, A14 (Nov. 2010), A14.
- [223] J. Bovy et al. “Think Outside the Color Box: Probabilistic Target Selection and the SDSS-XDQSO Quasar Targeting Catalog”. In: *Astrophys. J.* 729, 141 (Mar. 2011), p. 141. arXiv: 1011.6392 [astro-ph.CO].
- [224] N. P. Ross et al. “The SDSS-III Baryon Oscillation Spectroscopic Survey: Quasar Target Selection for Data Release Nine”. In: *Astrophys. J. Supp.* 199, 3 (Mar. 2012), p. 3. arXiv: 1105.0606 [astro-ph.CO].
- [225] A. Font-Ribera et al. “The large-scale quasar-Lyman α forest cross-correlation from BOSS”. In: *J. Cosmology Astropart. Phys.* 5, 018 (May 2013), p. 18. arXiv: 1303.1937 [astro-ph.CO].
- [226] N. P. Ross et al. “Clustering of Low-redshift ($z \leq 2.2$) Quasars from the Sloan Digital Sky Survey”. In: *Astrophys. J.* 697 (June 2009), pp. 1634–1655. arXiv: 0903.3230 [astro-ph.CO].
- [227] M. White et al. “The clustering of intermediate-redshift quasars as measured by the Baryon Oscillation Spectroscopic Survey”. In: *Mon. Not. R. Astron. Soc.* 424 (Aug. 2012), pp. 933–950. arXiv: 1203.5306 [astro-ph.CO].

- [228] A. Slosar et al. “The Lyman- α forest in three dimensions: measurements of large scale flux correlations from BOSS 1st-year data”. In: *J. Cosmology Astropart. Phys.* 9, 001 (Sept. 2011), p. 1. arXiv: 1104.5244 [astro-ph.CO].
- [229] K. B. Schmidt et al. “Selecting Quasars by Their Intrinsic Variability”. In: *Astrophys. J.* 714 (May 2010), pp. 1194–1208. arXiv: 1002.2642 [astro-ph.CO].
- [230] G. G. Williams et al. “90prime: a prime focus imager for the Steward Observatory 90-in. telescope”. In: *Ground-based Instrumentation for Astronomy*. Ed. by A. F. M. Moorwood and M. Iye. Vol. 5492. Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series. Sept. 2004, pp. 787–798.
- [231] E. L. Wright et al. “The Wide-field Infrared Survey Explorer (WISE): Mission Description and Initial On-orbit Performance”. In: *Astron. J.* 140, 1868 (Dec. 2010), pp. 1868–1881. arXiv: 1008.0031 [astro-ph.IM].
- [232] D. Lang. “unWISE: Unblurred Coadds of the WISE Imaging”. In: *Astron. J.* 147, 108 (May 2014), p. 108. arXiv: 1405.0308 [astro-ph.IM].
- [233] K. N. Abazajian et al. “The Seventh Data Release of the Sloan Digital Sky Survey”. In: *Astrophys. J. Supp.* 182, 543 (June 2009), pp. 543–558. arXiv: 0812.0649.
- [234] PanSTARRS. *PanSTARRS Survey Website*. <http://pan-starrs.ifa.hawaii.edu/public>. 2010.
- [235] N. M. Law et al. “The Palomar Transient Factory: System Overview, Performance, and First Results”. In: *Proc. Astron. Soc. Pacific* 121 (Dec. 2009), pp. 1395–1408. arXiv: 0906.5350 [astro-ph.IM].
- [236] SCUSS. *South Galactic Cap U-Band Sky Survey*. <http://batc.bao.ac.cn/Uband>. 2010.
- [237] D. Lang. “ ” in preparation. 2014.
- [238] R. H. Hardin, N. J. A. Sloane, and W. D. Smith. *Tables of Spherical Codes with Icosahedral Symmetry*. Florham Park: AT&T Shannon Lab., 2001.